

Effect of different non-ionic surfactants on the biodegradation of PAHs by diverse aerobic bacteria

L. Fernando Bautista^{a,*}, Raquel Sanz^a, M. Carmen Molina^b, Natalia González^b, David Sánchez^a

^a Department of Chemical and Environmental Technology, ESCET, Universidad Rey Juan Carlos, C/ Tulipán s/n, 28933 Móstoles, Madrid, Spain

^b Department of Biology and Geology, ESCET, Universidad Rey Juan Carlos, C/ Tulipán s/n, 28933 Móstoles, Madrid, Spain

ARTICLE INFO

Article history:

Received 6 January 2009

Received in revised form

24 June 2009

Accepted 25 June 2009

Available online 17 July 2009

Keywords:

Polycyclic aromatic hydrocarbons

Surfactants

Biodegradation

Aerobic bacteria

ABSTRACT

The aim of this work was to evaluate the effect of several non-ionic surfactants (Tween-80, Triton X-100 and Tergitol NP-10) on the ability of different bacteria (*Enterobacter* sp., *Pseudomonas* sp. and *Stenotrophomonas* sp.) to degrade polycyclic aromatic hydrocarbons (PAHs). Bacterial cultures were performed at 25 °C in an orbital shaker under dark conditions in BHB medium containing 1% of surfactant and 500 mg l⁻¹ of each PAH. Experiments performed with Tween-80 showed the highest cell density values and maximum specific growth rate because this surfactant was used as a carbon source by all bacteria. High degree of PAHs degradation (>90%) was reached in 15 days in all experiments. Toxicity increased at early times using Tween-80 but decreased to low levels in a short time after the firsts 24 h. On the other hand, Triton X-100 and Tergitol NP-10 were not biodegraded and toxicity kept constant along time. However, PAHs-degradation rate was higher, especially by the action of *Enterobacter* sp. with Tween-80 or Triton X-100. Control experiments performed without surfactant showed a significant decrease in biomass growth rate with a subsequent loss of biodegradation activity likely due to a reduced solubility and bioavailability of PAHs in absence of surfactant.

© 2009 Elsevier Ltd. All rights reserved.

1. Introduction

Polycyclic aromatic hydrocarbons (PAHs) are comprised of two or more fused aromatic rings. They are generally formed by incomplete combustion of fossil fuels and other organic material. They are known to be carcinogenic to humans and other organisms (Kim et al., 2001). PAHs should be undetectable in soil and groundwater under the drinking water standards and health advisory regulations of the US Environmental Protection Agency (USEPA), where 16 PAHs have been defined as priority pollutants and form a subset of “toxic pollutants” under the Clean Water Act including naphthalene, phenanthrene and anthracene (USEPA, 1972). Naphthalene is also considered a hazardous air pollutant (USEPA, 1994, 2004). These administrative regulations have encouraged scientists and engineers to identify the best remediation technologies from a variety of physical, chemical and biological methods that can be used to remove these compounds from polluted sites. Most low molecular weight PAHs are biodegradable in presence of a suitable microbial population. However, the efficiency of PAHs biodegradation is limited because these compounds have a very low aqueous solubility and vapour pressure (Luning Prak and Pritchard, 2002).

Surfactants may be useful for the bioremediation of sites polluted with PAHs since they enhance the solubility of hydrophobic compounds (Boonchan et al., 1998). Many studies have been conducted to enhance the biodegradation of PAHs using surfactants to increase their solubility by decreasing the interfacial surface tension between PAHs and the soil/water interphase. When surfactant concentration is above the critical micelle concentration (CMC), micelle aggregates provide an additional hydrophobic area in the central region of micelles enhancing the aqueous solubility of PAHs (Li and Chen, 2009). The addition of non-ionic surfactants as additives has a positive effect on PAHs biodegradation (Grimberg et al., 1996; Volkering et al., 1995). However, other studies have reported negligible (Ghosh et al., 1995) or even negative effects (Laha and Luthy, 1991) of surfactants. Possible reasons for this include the competition for substrate utilization and the toxicity of the surfactants toward PAHs-degrading bacteria (Liu et al., 2001). Therefore, PAHs-degradation processes involving surfactant utilization need to be optimized for each of the factors influencing biodegradation, including surfactants type and concentration, PAH specificity and the microorganisms present in the process (Jin et al., 2007).

The purpose of the present study was to examine the effect of the presence of non-ionic surfactants (Tween-80, Triton X-100 and Tergitol NP-10) on the biodegradation of low molecular weight PAHs (naphthalene, phenanthrene and anthracene) in aqueous media

* Corresponding author. Tel.: +34 91 488 8501; fax: +34 91 488 7068.

E-mail address: fernando.bautista@urjc.es (L. Fernando Bautista).

inoculated with one of each of the aerobic bacterial strains identified as *Pseudomonas* sp., *Enterobacter* sp. and *Stenotrophomonas* sp. or with a mixed culture comprised of these three species.

2. Material and methods

2.1. Chemicals and media

Naphthalene, phenanthrene and anthracene (all >99% purity) used in the degradation experiments were purchased from Sigma-Aldrich (Steinheim, Germany) and Fluka (Steinheim, Germany). Reagent grade dichloromethane was supplied by Scharlau Chemie (Barcelona, Spain). Bushnell Haas Broth medium (BHB) was purchased from Panreac (Barcelona, Spain). All surfactants (Tween-80, Triton X-100 and Tergitol NP-10) were purchased from Sigma-Aldrich (Steinheim, Germany). Relevant properties of these surfactants are shown in Table 1. All chemicals used for toxicity assays were at least reagent grade.

2.2. Microorganisms

Pseudomonas sp., *Enterobacter* sp. and *Stenotrophomonas* sp. were isolated and identified from a microbial consortium obtained from a soil chronically exposed to petroleum products from a petrochemical complex in Puertollano, Spain.

2.3. Biodegradation of PAHs

Biodegradation experiments were performed in 3 replicates. 1.5 ml of the culture were inoculated in 150 ml BHB medium containing 1% w/w of the corresponding surfactant. The resulting initial concentration for each surfactant was 15.9 mM for Tergitol NP-10, 17.1 mM for Triton X-100 and 8.1 mM for Tween-80. Each PAH was added to a concentration of 500 mg l⁻¹. Erlenmeyer flasks were incubated at 200 rpm and 25 °C during 30 days. Samples were withdrawn at different time intervals to monitor PAHs depletion and cell density. Control cultures without the addition of any surfactant were conducted for all bacterial strains. In addition, an abiotic control without bacterial inoculum was performed in order to evaluate PAH depletion due to physicochemical processes. Cell density was monitored by measuring the absorbance at 600 nm using a Cary-500 spectrophotometer (Varian, Palo Alto, CA, USA).

The depletion rate ($-r_i$) of each PAH (i) during the stationary phase of microbial growth was fitted to a first-order kinetic model (equation (1)) with satisfactory results.

$$-r_i = -\frac{dC_i}{dt} = k_{A,i} \cdot C_i + k_{B,i} \cdot C_i, \quad (1)$$

where C is the concentration of PAHs measured by HPLC analysis, k_A is the apparent first-order kinetic constant due to abiotic processes, k_B is the apparent first-order kinetic constant due to biological processes. In order to evaluate the effect of the surfactant on the biodegradation of each PAH by different microorganisms, a two-way analysis of variance (ANOVA) with two factors, type of microorganism (four levels: *Pseudomonas*, *Enterobacter*, *Stenotrophomonas*

and without inoculum) and type of surfactant (four levels: Tween-80, Triton X-100, Tergitol NP-10 and absence of surfactant), was performed for each PAH using the k_B values. In addition, a second analysis of variance with two factors, type of PAH (three levels: naphthalene, phenanthrene and anthracene) and type of microorganism (four levels: *Pseudomonas*, *Enterobacter*, *Stenotrophomonas* and all three together) was carried with the k_B values from those where Tween-80 was used as surfactant. The analyses were performed using the software Statistica 6.0 (StatSoft, Inc., Tulsa, OK, USA). The variances were checked for homogeneity by the Cochran's test and Student Newman Keuls (SNK) test was used to discriminate among different treatments after significant F -test.

2.4. HPLC analysis

Naphthalene, phenanthrene and anthracene concentrations in the culture media were performed using a reversed-phase C18 column (Luna C18(2), 7.5 cm × 4.6 mm I.D., 3 μm particle size. Phenomenex, Torrance, CA, USA) with its corresponding guard column. The HPLC system was a ProStar 230 (Varian, Palo Alto, CA, USA) equipped with a quaternary pump and photodiode array UV/Vis and fluorescence detectors. 10 μl of filtered cultivation media were injected and then eluted from the column at a flowrate of 0.8 ml min⁻¹ using an acetonitrile:water gradient program as follows: isocratic 60:40 (0–2 min), gradient to 75:25 (2–14 min), isocratic 75:25 (14–15 min), gradient to 100:0 (15–16 min). The concentration each PAH was calculated from a standard curve based on peak area at a wavelength of 254 nm.

2.5. Toxicity assays

The toxicity monitoring during PAHs degradation was carried out using the Microtox™ assay Greene et al. (1988). The assay was run with the bioluminescent bacterium *Vibrio fischeri*, following the protocol suggested by Microbics Corporation (1992). Toxicity was expressed as the percentage of the *V. fischeri* inhibition after 15 min of incubation with the assayed sample at 15 °C.

2.6. Total organic carbon and pH

Total organic carbon (TOC) analyses during biodegradation experiments were performed using a TOC-VCSN analyzer (Shimadzu, Japan). Samples were injected into the oven at 680 °C. The combustion was catalyzed by Pt/Al₂O₃ catalyst. The CO₂ produced was measured by NDIR method. Monitoring of pH during PAHs-biodegradation experiments was performed using a GPL21 pH-meter (Crison, Barcelona, Spain).

3. Results

3.1. Effects of different surfactants on the bacterial growth

The growth curves of the three bacterial strains during the PAHs-biodegradation experiments (Fig. 1) show the expected latent, exponential and stationary phases. Depending on the surfactant used, the

Table 1
Relevant properties of surfactants.

Surfactant	Molecular formula	Molecular Weight	CMC ^{a,b} (mM)	Density ^b (g ml) ⁻¹
Tween-80	C ₁₈ H ₃₇ (C ₆ H ₉ O ₅)(OC ₂ H ₄) ₂₀ OH	1309	0.012	1.06
Triton X-100	C ₈ H ₁₇ C ₆ H ₄ (OC ₂ H ₄) ₁₀ OH	625	0.232	1.07
Tergitol NP-10	C ₁₅ H ₂₄ O(OC ₂ H ₄) ₁₀ OH	652	0.069	1.04

^a CMC: critical micelle concentration.

^b at 298 K.




Fig. 1. Effect of different surfactants on the growth of PAHs-degrading bacteria (*Pseudomonas* sp. ■, *Enterobacter* sp. ●, *Stenotrophomonas* sp. ▲, and a mixture of the three above ◆) during PAHs-biodegradation experiments (error bars denote standard deviation of 3 replicates). Cultures were performed using Tween-80 (A), Triton X-100 (B) and Tergitol NP-10 (C), and without surfactant (D).

length of these phases was different. The end of the exponential phase was reached approximately after 48 h in experiments performed in presence of Tween-80 while 150 h were necessary for Triton X-100 and Tergitol NP-10. The stationary phase was usually completed after 15 days of incubation in all experiments. According to previous experiments, the three strains used in the present work were able to use Tween-80 as their sole carbon source. On the other hand, growth was not detected in presence of Triton X-100 and Tergitol NP-10 under the same conditions (data not shown).

Biodegradation experiments carried out with PAHs showed that the growth rates were greater in media using Tween-80 as the surfactant, while the lower growth was achieved in those cultures performed without surfactant. In addition, the highest growth (Fig. 1) and maximum specific growth rate (Fig. 2) was observed when a mixed culture of the three strains (*Pseudomonas* sp., *Enterobacter* sp. and *Stenotrophomonas* sp.) was employed.

3.2. Biodegradation of PAHs

Figures 3–5 show, respectively, the biodegradation of naphthalene, phenanthrene and anthracene in the medium during the biodegradation experiments for different surfactants and bacteria. The addition of any surfactant increased the solubility,




Fig. 2. Specific growth rate (μ) of PAHs-degrading bacteria in different surfactants (error bars denote standard deviation of 3 replicates). Experiments with Triton X-100 and Tergitol NP-10 were not performed for the three-bacterium consortium.




Fig. 3. Effect of different surfactants on the naphthalene depletion in abiotic control ∇ and in cultures inoculated with PAHs-degrading strains: *Pseudomonas* sp. \blacksquare , *Enterobacter* sp. \bullet , *Stenotrophomonas* sp. \blacktriangle individually or with a mixture of the three strains \blacklozenge (error bars denote standard deviation of 3 replicates). Cultures were performed using Tween-80 (A), Triton X-100 (B) and Tergitol NP-10 (C), and without surfactant (D).

approximately, by a factor of 100 for naphthalene and 1000 for phenanthrene and anthracene (solubility in water of naphthalene, phenanthrene and anthracene at 25 °C are 30 mg l⁻¹, 1.3 mg l⁻¹ and 0.073 mg l⁻¹, respectively).

Most of PAHs added especially phenanthrene and anthracene, were non soluble at the beginning of each experiment. However, during the exponential growth phase, as soluble PAHs were consumed, non-soluble PAHs were progressively dissolved. At the end of the exponential growth all PAHs had been dissolved (data not shown). Therefore, from that time, PAHs measurements in each culture corresponded to the whole amount of PAHs in the medium. In all cases, the values of the biodegradation rate constant (k_B) were calculated in the stationary growth phase, where the PAHs measurements were the true total concentration of PAHs.

Naphthalene was rapidly biodegraded and removed between 200 and 400 h depending on the surfactants and bacteria used (Fig. 3). Using surfactants, PAHs were almost completely depleted after 30 days. The rate of naphthalene biodegradation was higher than phenanthrene and anthracene for each of the surfactants tested (Table 2).

From the results of the abiotic control (Figures 3–5), it was observed that both phenanthrene and anthracene were not significantly depleted in absence of microorganisms ($k_A \approx 0$ for phenanthrene and anthracene). On the other hand, under the same conditions a progressive depletion of naphthalene was observed, but

the rate of naphthalene depletion in controls was significantly slower than that observed in presence of PAHs-degrading bacteria. The values of k_B are shown in Table 2. The ANOVA performed with k_B values for each PAH (Table 3) showed that, in all three cases, interactions between both factors (i.e., type of microorganism and type of surfactant) were clearly significant, as it can be seen from the statistical parameters for naphthalene ($F_{6,24} = 135.49$, $p < 0.001$, $n = 36$), phenanthrene ($F_{6,24} = 326.92$, $p < 0.001$, $n = 36$) and anthracene ($F_{6,24} = 70.62$, $p < 0.001$, $n = 36$). In all experiments, the presence of surfactant significantly improved the biodegradation process. For naphthalene biodegradation (Table 2), the best treatment was achieved with Tween-80 and the bacterial consortium, followed by *Enterobacter* sp. (SNK $p < 0.05$). Using this surfactant the *Pseudomonas* sp. strain showed the poorest naphthalene biodegradation capability among all three strains (SNK $p < 0.05$). For phenanthrene, Tween-80 and *Enterobacter* sp. showed the best results. However, *Pseudomonas* sp. strain yielded better results than *Stenotrophomonas* sp. strain for both Tween-80 and Triton X-10 (SNK $p < 0.05$). For the biodegradation of anthracene, *Enterobacter* sp. was again the best strain, although no significant differences were observed when using Tween-80 and Triton X-10 (SNK $p < 0.05$). With respect to the use of Tergitol NP-10, the best biodegradation rate for phenanthrene and anthracene was achieved by *Pseudomonas* sp., while the efficiency of *Enterobacter* sp. was significantly reduced compared to experiments where Tween-80




Fig. 4. Effect of different surfactants on the phenanthrene depletion in abiotic control ∇ and in cultures inoculated with PAHs-degrading strains: *Pseudomonas* sp. \blacksquare , *Enterobacter* sp. \bullet , *Stenotrophomonas* sp. \blacktriangle individually or with a mixture of the three strains \blacklozenge (error bars denote standard deviation of 3 replicates). Cultures were performed using Tween-80 (A), Triton X-100 (B) and Tergitol NP-10 (C), and without surfactant (D).

was used (SNK $p < 0.05$). These data were consistent with the results from the cell growth measurements.

3.3. Toxicity

The toxicity monitoring during the PAHs-degradation process (Fig. 6) showed a small increase during the exponential growth phase. This effect was likely due to metabolites formed and secreted into the medium. However, an overall decreasing trend was observed when Tween-80 was used as surfactant. In the case of other surfactants, the toxicity was constant during the time course of the experiment. This was due to the fact that Tween-80 is biodegradable and it is used as carbon source by the microorganisms while Tergitol NP-10 and Triton X-100 are not metabolised by the PAHs-degrading bacteria, showing that these surfactants were toxic for *V. fischeri*.

For the experiments performed with Tween-80 and the consortium of three bacteria, the toxicity decreased until values below 50%. These data show that the addition of some surfactants may decrease the toxicity of the medium.

3.4. TOC and pH

For a complete description of the biodegradation process, monitoring of pH and TOC from the supernatant were performed

(Figs. 7 and 8, respectively) during the 30 days of the incubation of each experiment. TOC evolution was similar in all experiments showing that at early times of the incubation, TOC increased reaching a maximum and then, sharply decreased along the stationary phase. At the end of the experiment TOC values slightly increased probably due to the release of compounds in the cultivation media during the death phase. In the experiment in absence of surfactant low TOC values in solution were reported and their evolution was slower than in experiments performed in presence of surfactants.

In all cases, pH increased during the exponential phase. For the experiments with Tween-80 as surfactants pH increased from 7.0 to approximately 9.0. While, with the others surfactants pH reached values of 7.8. At the end of experiments pH decreased slightly.

4. Discussion

Data show that the addition of surfactants affect positively PAHs biodegradation, increasing the solubility of PAHs and allowing their progressive biodegradation up to concentration levels lower than the initial values. This effect was clearly observed considering the low solubility and biodegradation degree of PAHs when surfactant was not added. The abiotic depletion of naphthalene in presence of 1% w/w Tween-80 ($k_A = 2.2 \times 10^{-3} \pm 6.0 \times 10^{-5} \text{ h}^{-1}$), was not




Fig. 5. Effect of different surfactants on the anthracene depletion in abiotic control ∇ and in cultures inoculated with PAHs-degrading strains: *Pseudomonas* sp. \blacksquare , *Enterobacter* sp. \bullet , *Stenotrophomonas* sp. \blacktriangle individually or with a mixture of the three strains \blacklozenge (error bars denote standard deviation of 3 replicates). Cultures were performed using Tween-80 (A), Triton X-100 (B) and Tergitol NP-10 (C), and without surfactant (D).

observed for both three-ring PAHs ($k_A \approx 0$), and it was likely due to volatilization caused by a higher vapour pressure of naphthalene (11.6 Pa) compared to that of phenanthrene (0.13 Pa) and anthracene (0.08 Pa). However, in presence of PAHs-degrading strains, naphthalene depletion was significantly faster than that in the

abiotic control experiment demonstrating that degradation of naphthalene was mainly due to biological processes.

Of the three surfactants studied, Tween-80 was the one that caused the higher level of PAHs degradation. Moreover, the progressive decrease in toxicity and the larger bacterial growth

Table 2
Apparent first-order biotic degradation rate constant (k_B) for each PAH with different microorganisms and surfactants.

Microorganism	Surfactant	k_B (h^{-1}) ^a		
		Naphthalene	Phenanthrene	Anthracene
<i>Pseudomonas</i> sp.	Tween-80	$7.9 \times 10^{-3} \pm 6.4 \times 10^{-4}$	$8.2 \times 10^{-3} \pm 2.0 \times 10^{-4}$	$3.3 \times 10^{-3} \pm 6.7 \times 10^{-4}$
	Triton X-100	$9.5 \times 10^{-3} \pm 4.4 \times 10^{-4}$	$8.6 \times 10^{-3} \pm 2.5 \times 10^{-4}$	$6.4 \times 10^{-3} \pm 2.3 \times 10^{-4}$
	Tergitol NP-10	$1.1 \times 10^{-2} \pm 3.3 \times 10^{-4}$	$5.3 \times 10^{-3} \pm 3.2 \times 10^{-4}$	$5.7 \times 10^{-3} \pm 3.1 \times 10^{-4}$
	No surfactant	$2.4 \times 10^{-3} \pm 1.8 \times 10^{-4}$	$1.0 \times 10^{-3} \pm 1.3 \times 10^{-5}$	$4.7 \times 10^{-4} \pm 7.3 \times 10^{-6}$
<i>Enterobacter</i> sp.	Tween-80	$1.7 \times 10^{-2} \pm 2.2 \times 10^{-4}$	$1.2 \times 10^{-2} \pm 7.4 \times 10^{-5}$	$7.0 \times 10^{-3} \pm 5.3 \times 10^{-4}$
	Triton X-100	$1.2 \times 10^{-2} \pm 3.9 \times 10^{-5}$	$9.8 \times 10^{-3} \pm 3.7 \times 10^{-5}$	$7.3 \times 10^{-3} \pm 1.4 \times 10^{-4}$
	Tergitol NP-10	$1.1 \times 10^{-2} \pm 6.6 \times 10^{-5}$	$3.3 \times 10^{-3} \pm 1.3 \times 10^{-4}$	$4.4 \times 10^{-3} \pm 1.2 \times 10^{-4}$
	No surfactant	$4.0 \times 10^{-3} \pm 3.7 \times 10^{-4}$	$5.2 \times 10^{-4} \pm 1.2 \times 10^{-5}$	$5.5 \times 10^{-4} \pm 3.4 \times 10^{-5}$
<i>Stenotrophomonas</i> sp.	Tween-80	$1.5 \times 10^{-2} \pm 7.4 \times 10^{-4}$	$7.4 \times 10^{-3} \pm 1.1 \times 10^{-4}$	$5.2 \times 10^{-3} \pm 4.7 \times 10^{-4}$
	Triton X-100	$1.3 \times 10^{-2} \pm 1.7 \times 10^{-5}$	$5.3 \times 10^{-3} \pm 1.6 \times 10^{-4}$	$3.1 \times 10^{-3} \pm 1.0 \times 10^{-4}$
	Tergitol NP-10	$1.1 \times 10^{-2} \pm 3.3 \times 10^{-4}$	$2.0 \times 10^{-3} \pm 1.2 \times 10^{-4}$	$3.1 \times 10^{-3} \pm 1.2 \times 10^{-4}$
	No surfactant	$9.8 \times 10^{-4} \pm 3.2 \times 10^{-5}$	$7.8 \times 10^{-4} \pm 4.7 \times 10^{-5}$	$5.2 \times 10^{-4} \pm 1.5 \times 10^{-5}$
3 bacteria above	Tween-80	$2.0 \times 10^{-2} \pm 3.9 \times 10^{-5}$	$9.5 \times 10^{-3} \pm 1.5 \times 10^{-4}$	$6.7 \times 10^{-3} \pm 5.5 \times 10^{-6}$
	No surfactant	$4.6 \times 10^{-3} \pm 1.9 \times 10^{-4}$	$8.7 \times 10^{-4} \pm 4.0 \times 10^{-5}$	$5.6 \times 10^{-4} \pm 4.1 \times 10^{-5}$

^a \pm values correspond to standard deviation of 3 replicates.

Table 3
Analysis of variance (ANOVA) summary for the biotic degradation rate constant (k_B).

	Source	Sum of squares	D.F. ^a	F-value	p-value
Naphthalene ^f	Intercept	0.003395	1	25529.64	<0.0001
	Surfactant ^b	0.000655	3	1642.96	<0.0001
	Strain ^c	0.000073	2	274.71	<0.0001
	Interaction	0.000108	6	135.49	<0.0001
	Error	0.000003	24		
Phenanthrene ^g	Intercept	0.001036	1	45705.99	<0.0001
	Surfactant ^b	0.000412	3	6053.49	<0.0001
	Strain ^c	0.000041	2	914.81	<0.0001
	Interaction	0.000044	6	326.92	<0.0001
	Error	0.000001	24		
Anthracene ^h	Intercept	0.000553	1	5725.337	<0.0001
	Surfactant ^b	0.000147	3	506.049	<0.0001
	Strain ^c	0.000020	2	105.797	<0.0001
	Interaction	0.000041	6	70.616	<0.0001
	Error	0.000002	24		
Tween-80 ⁱ	Intercept	0.003605	1	21129.93	<0.0001
	PAHs ^d	0.000561	2	1645.28	<0.0001
	Strain ^e	0.000198	3	387.24	<0.0001
	Interaction	0.000114	6	111.03	<0.0001
	Error	0.000004	24		

^a Degrees of freedom.

^b Surfactant: Tween-80, Triton X-100, Tergitol NP-10 and without surfactant.

^c Strain: *Pseudomonas* sp., *Enterobacter* sp. and *Stenotrophomonas* sp.

^d PAHs: naphthalene, phenanthrene and anthracene.

^e Strain: *Pseudomonas* sp., *Enterobacter* sp., *Stenotrophomonas* sp and without inoculation.

^f Effect of the type of strain and surfactant on the biodegradation of naphthalene.

^g Effect of the type of strain and surfactant on the biodegradation of phenanthrene.

^h Effect of the type of strain and surfactant on the biodegradation of anthracene.

ⁱ Effect of the type of PAH and strain on the biodegradation process using Tween-80 as surfactant.

seem to favour a significant reduction in naphthalene, phenanthrene and anthracene. For Triton X-100 and Tergitol NP-10 the results obtained were favourable and similar in both cases. However, the degree and rates of biodegradation were lower compared to the results using Tween-80. For both Triton X-100 and Tergitol NP-10 complete degradation of phenanthrene and anthracene was not achieved after 30 days of incubation.

Some studies have suggested positive effects similar to that reported in the present work both for Tween-80 and Triton X-100 (Kim et al., 2001; Yang et al., 2003; Cheng and Wong, 2006) and for Tergitol NP-10 (Volkering et al., 1995; Grimberg et al., 1996). However, other authors (Laha and Luthy, 1991, 1992; Yuan et al., 2000) reported inhibitory effects of the bacterial degradation activity when the surfactants were added in concentrations over the CMC, likely due to an excess of micelles of the pollutant and a reduction of the bioavailability (Mulligan et al., 2001). However, this effect was not observed in the present study, since the addition of surfactants high above CMC (Table 1) produced positive results.

The higher bacterial growth and PAHs-biodegradation rate observed in presence of Tween-80 during PAHs-biodegradation experiments could be due to the fact that this surfactant was biodegradable and can be used as an additional carbon source for the microorganisms (Franzetti et al., 2006; Boonchan et al., 1998; Yoshimura, 1986; Swisher, 1987), although the surfactant biodegradation could compete for the substrates (Tiehm, 1994).

The growth of *Pseudomonas* sp., *Enterobacter* sp. and *Stenotrophomonas* sp. strains with PAHs as carbon source indicates that these microorganisms are potential PAHs' biodegraders. The results showed that in most cases growth rate of *Enterobacter* sp. and its PAHs-biodegradation rate were higher than for the other bacteria. Furthermore, the rates of degradation and growth observed for




Fig. 6. Toxicity of culture media during PAHs-biodegradation experiments using different surfactants Tween-80 (*Pseudomonas* sp. ■, *Enterobacter* sp. ●, *Stenotrophomonas* sp. ▲ a mixture of the three strains ◆, abiotic control ▽); Triton X-100 (*Pseudomonas* sp. □); and Tergitol NP-10 (*Pseudomonas* sp. ○).

Pseudomonas sp. and *Stenotrophomonas* sp. were also significant. Many authors have studied the genes involved in the PAHs-degradation metabolism for naphthalene (Dunn and Gunsalus, 1973; Kuhm et al., 1991; White et al., 1997; Filonov et al., 2006), phenanthrene (Balashova et al., 1999; Puntus et al., 2008) and anthracene (Jacques et al., 2005) or other aromatic hydrocarbon with higher molecular weight (Boonchan et al., 1998; Juhasz et al., 2000). Anyway, the highest levels of growth and degradation of PAHs were obtained when the three bacteria worked together. Probably this fact is due to the existence of a microbial succession during the incubation.

The PAHs-degrading efficiency depends on the molecular weight and the number of aromatic rings (Warner and Peters, 2005), since, the solubility in water, volatility and persistence of PAHs decrease when molecular weight and the number of aromatic rings decrease (Juhasz and Naidu, 2000; Kanaly and Harayama, 2000; Dimitriou-Christidis et al., 2007). The results showed that the evolution of the PAHs during the biodegradation process followed the order of solubility of these pollutants: naphthalene > phenanthrene > anthracene (Figures 3–5). The depletion of both three-ring hydrocarbons was not associated to any abiotic processes since depletion of phenanthrene and anthracene was not observed from the abiotic control experiments. In the case of naphthalene, the depletion measured in this abiotic control was likely due to volatilization because of its higher vapour pressure (11.6 Pa at 298 K) than phenanthrene and anthracene (0.13 Pa and 0.08 Pa at 298 K, respectively). However, the overall rate of naphthalene depletion was higher than that observed in control experiments, so that the biological contribution was significant. The large increase in the degradation rate (eq. (1)) of each PAH in presence of surfactant was not only due to the large increase in PAHs solubility, but to a concomitant increase in the value of the kinetic constant (Table 2). The significant interactions between type of microorganism and type of surfactant showed by the ANOVA performed for each PAH prove that the choice of the best surfactant to increase the biodegradation efficiency depends on the microorganism or consortium used as well as the nature of the hydrocarbon to be removed.




Fig. 7. Total organic carbon (TOC) during PAHs-biodegradation experiments by diverse PAHs-degrading bacteria (*Pseudomonas* sp. ■, *Enterobacter* sp. ●, *Stenotrophomonas* sp. ▲, a mixture of the three strains ◆). Cultures were performed using Tween-80 (A), Triton X-100 (B) and Tergitol NP-10 (C), and without surfactant (D).

Most studies report a depletion of target components without considering the toxicity during the process. Fig. 6 shows that at the beginning of each experiment toxicity was constant or even slightly increased as a consequence of the potential accumulation of intermediate products formed (Grifoll et al., 1995). Biodegradation experiments using *Pseudomonas* sp. with Triton X-100 or Tergitol NP-10 did not show a significant decrease in toxicity along the whole course of both experiments (Fig. 6). Although toxic effects produced by chemical surfactants on bacterial communities have been reported (Kim et al., 2001), in this work, this effect was not observed, at least to a large extent, since *Pseudomonas* sp. showed capability to degrade PAHs almost completely. This means that these two surfactants were not toxic for *Pseudomonas* sp. but induced toxicity for *V. fischeri*, i.e., the bacteria used for the toxicity assay. However, when Tween-80 was used as surfactant, a significant decrease in toxicity was achieved (Fig. 6) since, not only PAHs were depleted but, all three strains are capable of metabolizing the surfactant. In the absence of surfactant a lower decrease in toxicity was shown.

The small decrease in TOC (Fig. 7) confirmed the PAHs biodegradation. The partial mineralization of PAHs and intermediate products to CO₂ could explain this trend. Nevertheless the results showed two increases in TOC levels during the biodegradation process. The first one, from the beginning of the experiment, could be caused by the formation of intermediate degradation products

and the dissolution of PAHs precipitated in the media along the exponential growth phase. In the case of the biodegradation experiments using Tween-80 as surfactant, that effect was even larger (Fig. 7A) since Tween-80 was biodegraded by all three strains used. At the end of the growth phase a second increase in TOC took place, most likely due to the release of intracellular matter during the death phase. In absence of surfactant, the low solubility of PAHs produced very low values of TOC in aqueous solution.

An increase in pH was observed in all experiments, probably due to intermediate products formed. At the beginning of each experiment metabolites bearing hydroxyl groups increase the pH of the culture media (Bossert and Bartha, 1984; Juhasz and Naidu, 2000; Habe and Omori, 2003; Luan et al., 2006; Puntus et al., 2008). The higher pH increase when Tween-80 was used is likely due to the excretion of surfactant metabolites in the culture medium (Dibble and Bartha, 1979).

In conclusion, this study showed the positive effect of non-ionic surfactants on the PAHs-biodegradation process due to a remarkable increase in both solubility and bioavailability of these hydrocarbons, yielding higher biodegradation rates in presence of surfactants. The highest values of PAHs degradation and bacterial growth rate were obtained when Tween-80 was used as surfactant. Furthermore, this surfactant was used as a growth substrate by the bacteria. The strains used in this work completely degraded two and three-ring hydrocarbons. The best results were obtained with




Fig. 8. pH during PAHs-biodegradation experiments by diverse PAHs-degrading bacteria (*Pseudomonas* sp. ■, *Enterobacter* sp. ●, *Stenotrophomonas* sp. ▲, a mixture of the three strains ◆). Cultures were performed using Tween-80 (A), Triton X-100 (B) and Tergitol NP-10 (C), and without surfactant (D).

the use of a consortium composed of the three bacteria for the biodegradation of naphthalene and *Enterobacter* sp. for phenanthrene and anthracene.

Acknowledgements

Authors gratefully acknowledge the financial support from the Spanish Ministry of Environment (Research Project 13/2006/2-1.1) and Fundación Alfonso Martín Escudero.

References

- Balashova, N.V., Kosheleva, I.A., Golovchenko, N.P., Boronin, A.M., 1999. Phenanthrene metabolism by *Pseudomonas* and *Burkholderia* strains. *Process Biochemistry* 35, 291–296.
- Boonchan, S., Britz, M.L., Stanley, G.A., 1998. Surfactant-enhanced biodegradation of high molecular weight polycyclic aromatic hydrocarbons by *Stenotrophomonas maltophilia*. *Biotechnology and Bioengineering* 59, 482–494.
- Bossert, I., Bartha, R., 1984. The fate of petroleum in soils ecosystems. In: Atlas, R.M. (Ed.), *Petroleum Microbiology*. Macmillan Publishing Co., New York, pp. 435–489.
- Cheng, K.Y., Wong, J.W.C., 2006. Effect of synthetic surfactants on the solubilization and distribution of PAHs in water/soil-water systems. *Environmental Technology* 27, 835–844.
- Dibble, J.R., Bartha, R., 1979. Effect of environmental parameters on the biodegradation of oil sludge. *Applied and Environmental Microbiology* 37, 729–739.
- Dimitriou-Christidis, P., Autenrieth, R.L., McDonald, T.J., Desai, A.M., 2007. Measurement of biodegradability parameters for single unsubstituted and methylated polycyclic aromatic hydrocarbons in liquid bacterial suspensions. *Biotechnology and Bioengineering* 97, 922–932.
- Dunn, N.W., Gunsalus, I.C., 1973. Transmissible plasmid coding early enzymes of naphthalene oxidation in *Pseudomonas putida*. *Journal of Bacteriology* 114, 974–979.
- Filonov, A.E., Puntus, I.F., Karpov, A.V., Kosheleva, I.A., Akhmetov, L.I., Yonge, D., Petersen, J., Boronin, A.M., 2006. Assessment of naphthalene biodegradation efficiency provided by microorganisms of genera *Pseudomonas* and *Burkholderia* in soil model systems. *Journal of Chemical Technology and Biotechnology* 81, 216–224.
- Franzetti, A., Di Gennaro, P., Bevilacqua, A., Papacchini, M., Bestetti, G., 2006. Environmental features of two commercial surfactants widely used in soil remediation. *Chemosphere* 62, 1474–1480.
- Ghosh, M.M., Yeom, I.T.M., Shi, Z., Cox, C.D., Robinson, K.G., 1995. *Microbial Processes for Bioremediation*. Battelle Press, Columbus, OH, USA.
- Greene, J.C., Bartels, C.L., Warre-Hicks, W.J., Parkhurst, B.R., Linder, G.L., Peterson, S.A., Miller, W.E., 1988. *Protocols for Short Term Toxicity Screening of Hazardous Waste Sites*. In: EPA/600/3-88/029. US-EPA, Corvallis, OR, USA.
- Grifoll, M., Awlindonoc, S.A., Gatlin, C.V., Chapman, P.J., 1995. Action of a versatile fluorine-degrading bacterial isolate on polycyclic aromatic compounds. *Applied and Environmental Microbiology* 61, 3711–3723.
- Grimberg, S.J., Stringfellow, W.T., Aitken, M.D., 1996. Quantifying the biodegradation of phenanthrene by *Pseudomonas stutzeri* P16 in the presence of a nonionic surfactant. *Applied and Environmental Microbiology* 62, 2387–2392.
- Habe, H., Omori, T., 2003. Genetics of polycyclic aromatic hydrocarbon metabolism in diverse aerobic bacteria. *Bioscience Biotechnology and Biochemistry* 67, 225–243.
- Jacques, R.J., Santos, E.C., Bento, F.M., Perlaba, M.C., Selbach, P.A., Sá, E.L., Camargo, F.A., 2005. Anthracene biodegradation by *Pseudomonas* sp. isolated from a petrochemical sludge landfarming site. *International Biodeterioration and Biodegradation* 56, 143–150.
- Jin, D., Juang, X., Jing, X., Ou, Z., 2007. Effect of concentration, head group, and structure of surfactants on the degradation of phenanthrene. *Journal of Hazardous Materials* 144, 215–221.
- Juhász, A.L., Naidu, R., 2000. Bioremediation of high molecular weight polycyclic aromatic hydrocarbons: a review of the microbial degradation of benzo[a]pyrene. *International Biodeterioration and Biodegradation* 45, 57–88.

- Juhasz, A.L., Stanley, G.A., Britz, M.L., 2000. Microbial degradation and detoxification of high molecular weight polycyclic aromatic hydrocarbons by *Stenotrophomonas maltophilia* strain VUN 10,003. *Letters in Applied Microbiology* 30, 396–401.
- Kanally, R.A., Harayama, S., 2000. Biodegradation of high-molecular-weight polycyclic aromatic hydrocarbons by bacteria. *Journal of Bacteriology* 182, 2059–2067.
- Kim, I.S., Park, J.S., Kim, K.W., 2001. Enhanced biodegradation of polycyclic aromatic hydrocarbons using non-ionic surfactants in soil slurry. *Applied Geochemistry* 16, 1419–1428.
- Kuhm, A.E., Stolz, A., Knackmuss, H.J., 1991. Metabolism of naphthalene by the biphenyl-degrading bacterium *Pseudomonas paucimobilis* Q1. *Biodegradation* 2, 115–120.
- Laha, S., Luthy, R.G., 1991. Inhibition of phenanthrene mineralization by nonionic surfactants in soil–water systems. *Environmental Science and Technology* 25, 1920–1930.
- Laha, S., Luthy, R.G., 1992. Effects of nonionic surfactants on the solubilization and mineralization of phenanthrene in soil–water systems. *Biotechnology and Bioengineering* 40, 1367–1380.
- Li, J.L., Chen, B.H., 2009. Effect of non-ionic surfactants on biodegradation of phenanthrene by a marine bacteria of *Neptunomonas naphthovorans*. *Journal of Hazardous Materials* 162, 66–73.
- Liu, Y., Zhu, L., Shen, X., 2001. Polycyclic aromatic hydrocarbons (PAHs) in indoor and outdoor air of Hangzhou, China. *Environmental Science and Technology* 35, 840–844.
- Luan, T.G., Yub, K.S.H., Zhonga, Y., Zhou, H.W., Lana, C.Y., Tamb, N.F.Y., 2006. Study of metabolites from the degradation of polycyclic aromatic hydrocarbons (PAHs) by bacterial consortium enriched from mangrove sediments. *Chemosphere* 65, 2289–2296.
- Luning Prak, D.J., Pritchard, P.H., 2002. Solubilization of polycyclic aromatic hydrocarbon mixtures in micellar non-ionic surfactant solution. *Water Research* 36, 3463–3472.
- Microbics Corporation, 1992. *Microtox Manual*. In: *Condensed Protocols. A Toxicity Testing Handbook*, vol. III. Microbics Corp., Carlsbad, CA, USA.
- Mulligan, C.N., Young, R.N., Gibbs, B.F., 2001. Surfactant-enhanced remediation of contaminated soil: a review. *Engineering Geology* 60, 371–380.
- Puntus, I.F., Filonov, A.E., Akhmetov, L.I., Karpov, A.V., Boronin, A.M., 2008. Phenanthrene degradation by bacteria of the genera *Pseudomonas* and *Burkholderia* in model soil systems. *Microbiology* 77, 7–15.
- Swisher, R.D., 1987. *Surfactant Biodegradation*, second ed. Marcel Dekker, New York.
- Tiehm, A., 1994. Degradation of polycyclic aromatic hydrocarbons in the presence of synthetic surfactants. *Applied and Environmental Microbiology* 60, 258–263.
- USEPA, 1972. Clean Water Act (33 USC, 1251 et seq.). United States Environmental Protection Agency, Washington, DC. Available at: <http://www.epa.gov/lawsregs/laws/cwa.html> (accessed 15.06.2008).
- USEPA, 1994. National Emissions Standards for Hazardous Air Pollutants for Source Categories, Subpart F (40 CFR 63). United States Environmental Protection Agency, Washington, DC. Available at: <http://ecfr.gpoaccess.gov> (accessed 15.06.2008).
- USEPA, 2004. Clean Air Act (42 USC, 7401–7626). Available at: United States Environmental Protection Agency, Washington, DC. Available at: <http://www.epa.gov/air/caa/> (accessed 15.06.2008).
- Volkering, F., Breure, A.M., Van Anel, J.G., Rulkens, W.H., 1995. Influence of nonionic surfactants on bioavailability and biodegradation of polycyclic aromatic hydrocarbons. *Applied and Environmental Microbiology* 61, 1699–1705.
- Warmer, K.H., Peters, C.A., 2005. Polycyclic aromatic hydrocarbon biodegradation rates: a structure-based study. *Environmental Science and Technology* 39, 2571–2578.
- White, L.G., Bourbonnière, L., Greer, C.W., 1997. Biodegradation of petroleum hydrocarbons by psychrotrophic *Pseudomonas* strains possessing both alkane (alk) and naphthalene (nah) catabolic pathways. *Applied and Environmental Microbiology* 63, 3719–3723.
- Yang, J.G., Liu, X., Long, T., Yu, G., Peng, S., Zheng, L., 2003. Influence of nonionic surfactant on the solubilization and biodegradation of phenanthrene. *Journal of Environmental Sciences* 15, 859–862.
- Yoshimura, K., 1986. Biodegradation and fish toxicity of nonionic surfactants. *Journal of the American Oil Chemists' Society* 63, 1590–1596.
- Yuan, S.Y., Wei, S.H., Chang, B.V., 2000. Biodegradation of polycyclic aromatic hydrocarbons by a mixed culture. *Chemosphere* 41, 1463–1468.