

## Phylogenetic study of *Catapyrenium s. str.* (Verrucariaceae, lichen-forming Ascomycota) and related genus *Placidiopsis*

María Prieto<sup>1</sup>  
Isabel Martínez  
Gregorio Aragón  
Mónica A.G. Otálora

Área de Biodiversidad y Conservación, Universidad Rey  
Juan Carlos, C/ Tulipán s/n, 28933 Móstoles,  
Madrid, Spain

François Lutzoni

Department of Biology, Duke University, Box 90338,  
Durham, North Carolina 27708

**Abstract:** The current classification of what used to be called *Catapyrenium* comprises eight genera belonging to distinct lineages in the Verrucariaceae. Previous phylogenetic studies have shown that the redefined genus *Catapyrenium* (*Catapyrenium s. str.*) is monophyletic and sister of *Placidiopsis* within the *Staurothele* group, but this relationship was based on only two species from each genus. We conducted a phylogenetic study of *Catapyrenium* and *Placidiopsis* as currently delimited to evaluate the monophyly of each genus and infer infrageneric relationships. An initial family level phylogenetic analysis based on the nuLSU locus and implementing a backbone constraint tree (with both weighted maximum parsimony and bootstrap maximum likelihood approaches) was performed to infer phylogenetic placements of *Catapyrenium* and *Placidiopsis* taxa not included in previous molecular systematic studies. The results of this analysis were used to define the ingroup for a second phylogenetic analysis based on nuITS and nuLSU and centered on *Catapyrenium s. str.* and *Placidiopsis*. *Placidiopsis* was found to be monophyletic, whereas *Catapyrenium s. str.* was not. *Catapyrenium dactylinum* was found to be closely related to *Placopyrenium caeruleopulvinum* and *Placopyrenium stanfordii*, all of which were closely related to *Placocarpus schaeferi* and *Verrucula*. In addition we found genus *Placopyrenium* to be polyphyletic. The resulting trees confirmed that *Catapyrenium s. str.* (excluding *C. dactylinum*) and *Placidiopsis* constitute two sister monophyletic entities. The data do not support *Placidiopsis cinerascens* and *P. tenella* as two distinct species because no characters can be used to

distinguish them. Thus *P. tenella* is here reduced to synonymy with *P. cinerascens*.

**Key words:** ascospore septation, backbone constraint tree, *Catapyrenium s. str.*, lichens, phylogeny, *Placidiopsis*, *Placopyrenium s. str.*, synonymy, Verrucariaceae

### INTRODUCTION

*Catapyrenium* Flotow is a squamulose genus with perithecial ascomata belonging to the Verrucariaceae (Verrucariales, Ascomycota). The genus was split into eight genera (Harada 1993, Breuss 1996a), based on combinations of characters such as the type of pycnidium, ascus shape and arrangement of the ascospores, thallus anatomy and morphology (structure of the upper cortex and type of anchoring organs) and presence or absence of an involucrellum. Under this classification the eight genera of *Catapyrenium s.l.* are *Anthracoarpon* Breuss, *Catapyrenium* Flot. (*Catapyrenium s. str.*), *Clavascidium* Breuss, *Heteroplacidium* Breuss, *Involucropyrenium* Breuss, *Neocatapyrenium* H. Harada, *Placidium* A. Massal. and *Scleropyrenium* H. Harada.

*Catapyrenium s. str.* comprises eight squamulose species with a thin paraplectenchymatous upper cortex poorly delimited from the algal layer (called pseudocortex in Gueidan et al. 2007), perithecial ascomata, clavate asci with biseriolate ascospores and *Dermatocarpon*-type pycnidia (the latter based on *Catapyrenium dactylinum*, a recently described species) (Breuss 1996a, 2000, 2002a). They inhabit soil, detritus, mosses and bark (Breuss 2001, 2002a) and are distributed worldwide in arid, semi-arid and arctic-alpine regions.

One of the main features distinguishing *Catapyrenium s. str.* from other genera formerly classified in *Catapyrenium s.l.* is the type of upper cortex (*cinereum*-type sensu Breuss 1990, 1996a). In this genus the upper cortex is thin (less than 30 µm), not distinctly differentiated from the medulla and with small cells. *Anthracoarpon*, *Clavascidium*, *Heteroplacidium*, *Neocatapyrenium* and *Placidium* develop a well defined eucortex, clearly differentiated from the medulla, and with larger cells (*lachneum*-type sensu Breuss 1990, 1996a). *Scleropyrenium* is the only genus that has a pachydermatous upper cortex (Harada 1993). Another important character that segregates this complex is a *Dermatocarpon*-type pycnidium,

Submitted 6 Jul 2009; accepted for publication 3 Sep 2009.

<sup>1</sup> Corresponding author. E-mail: maria.prieto@urjc.es

present in *Catapyrenium s. str.* (based only on *Catapyrenium dactylinum*), *Clavascidium*, *Heteroplacidium* and *Placidium*, while *Anthracoarpon*, *Neocatapyrenium* and *Scleropyrenium* have *Endocarpon*-type pycnidia (for explanations and pictures of the pycnidia types see Janex-Favre and Wagner 1986, Ménard and Roux 1995, Gueidan et al. 2007, Prieto et al. 2008). Members of genus *Involucropyrenium* share the same type of upper cortex with *Catapyrenium*; however the former has perithecia situated between the squamules, which have an involucrellum. The type of pycnidia has been described as *Endocarpon*-type based on one recently described species of *Involucropyrenium* (Roux 2005).

Other species not closely related to the *Catapyrenium s.l.* complex have been described or combined as *Catapyrenium* (i.e. *Catapyrenium caeruleopulvinum* and *Dermatocarpon zahlbruckneri*) and later subsumed within *Placopyrenium* Breuss by Breuss (2002c, 2009). *Placopyrenium* comprises species with crustose areolate to placodioid or subsquamulose thalli, some of them being parasitic species. The ascospores are colorless, simple to uni- or triseptate with pycnidia of *Dermatocarpon*-type (Breuss 2002c, 2009). Unlike *Catapyrenium s. str.*, *Placopyrenium* includes placodioid thalli and areoles fixed to the substrate by constricted bases or stipe-like holdfasts (TABLE I).

While *Placidiosis* Beltramini is very similar to *Catapyrenium s. str.*, sharing the same type of upper cortex, it differs in having septate ascospores and lacking pycnidia (TABLE I). This genus consists of 14 species worldwide (Breuss 1996b, 2002b), inhabiting soil, detritus, mosses and algal or lichen crusts. Most *Placidiosis* species are easily identified, except for *P. cinerascens* and *P. tenella*, which are more difficult to distinguish from each other. Both species were described by Nylander in the same year (1853a, b) on the basis of different ascospore lengths. However more than a century later Breuss (1994, 1996b) and Nimis and Martellos (2004) showed that by including more specimens the range of measurements for this character overlapped with the result that the presence of a small lid-like involucrellum in *P. tenella* became the main trait used to distinguish it from *P. cinerascens*.

Although ascospore septation is the character used to distinguish *Placidiosis* from *Catapyrenium s. str.*, molecular studies showed that in other groups of Ascomycota (Staiger 2002, Miller and Hundorf 2004) ascospore septation was not a reliable character for delineating monophyletic groups at genus and higher taxonomical ranks. Within the Verrucariaceae Savić et al. (2008) pointed out that the *Thelidium* group contained a mixture of species with nonseptate, transversely septate and muriform ascospores, due

TABLE I. Main diagnostic characters for distinguishing between genera included in *Catapyrenium s.l.*, *Placidiosis* and *Placopyrenium s.l.*

Genus	Growth form	Life strategy	Upper cortex	Attaching organ	Ascospore	Pycnidia type
<i>Anthracoarpon</i>	squamulose	nonparasitic	eucortex ( <i>lachneum</i> -type)	rhizohyphae and rhizines	simple	<i>Endocarpon</i>
<i>Catapyrenium s.str.</i>	squamulose	nonparasitic	pseudocortex ( <i>cinereum</i> -type)	rhizohyphae	simple	<i>Dermatocarpon</i> <sup>a</sup>
<i>Clavascidium</i>	squamulose	nonparasitic	eucortex ( <i>lachneum</i> -type)	rhizohyphae and rhizines	simple	<i>Dermatocarpon</i>
<i>Heteroplacidium</i>	crustose areolate to squamulose	nonparasitic	eucortex ( <i>lachneum</i> -type)	rhizohyphae and rhizines	simple	<i>Dermatocarpon</i>
<i>Involucropyrenium</i>	squamulose	nonparasitic	pseudocortex ( <i>cinereum</i> -type)	rhizohyphae	simple	<i>Endocarpon</i>
<i>Neocatapyrenium</i>	squamulose	nonparasitic	eucortex ( <i>lachneum</i> -type)	rhizohyphae and rhizines	simple	<i>Endocarpon</i>
<i>Placidiosis</i>	squamulose	nonparasitic	pseudocortex ( <i>cinereum</i> -type)	rhizohyphae and rhizines	uniseptate	—
<i>Placidium</i>	squamulose	nonparasitic	eucortex ( <i>lachneum</i> -type)	rhizohyphae and rhizines	simple	<i>Dermatocarpon</i>
<i>Placopyrenium</i>	crustose areolate to placodioid or subsquamulose	parasitic or not	pseudocortex	rhizines or stipes	simple to 1(–3)-septate	<i>Dermatocarpon</i>
<i>Scleropyrenium</i>	squamulose	nonparasitic	eucortex (pachydermatous)	rhizohyphae and rhizines	simple	<i>Endocarpon</i>

<sup>a</sup>Data for pycnidia type is based on *Catapyrenium dactylinum*.

to the homoplasious nature of these traits. There are also examples at the infrageneric level where ascospore septation has been shown to be strongly homoplasious (Ihlen and Ekman 2002, Naesborg et al. 2007). Otálora et al. (2010) revealed that ascospore septation is homoplasious even at the species level within the Collemataceae. Gueidan et al. (2007) showed nonseptate ascospores to be a symplesiomorphic character state retained in several distinct lineages across the Verrucariaceae, together with convergent and parallel evolution of various types of ascospore septation occurring at different rates among clades, thus rendering more difficult our understanding of the evolution of ascospore septation within this family.

Breuss (1983, 1996b) and Breuss and Hansen (1988) noted that, because the only character state separating *Catapyrenium s. str.* from *Placidiopsis* was the presence of a transverse septum in the ascospores of *Placidiopsis*, the separation of these two genera was in need of reappraisal. Specimens of *Catapyrenium s. str.* with uniseptate ascospores, as well as some *Placidiopsis* specimens with nonseptate ascospores, have been observed by the first author and were reported by Gueidan et al. (2007). Breuss and Hansen (1988) pointed out that ascospore septation in *Placidiopsis* ("true" septa) was distinct from *Catapyrenium s. str.* (pseudosepta).

On the other hand the ascus apex of *Placidiopsis* has been reported to differ from that of *Catapyrenium s. str.* in having a slight thickening in the apical ascus wall (tholus) showing an ocular chamber (i.e. a dome-shaped indentation of the lower edge of the tholus) (Harris 1979, Breuss 2002b, Nimis and Martellos 2004). However this feature has not been observed in all species of *Placidiopsis*, being visible only at a certain developmental stage (Breuss pers comm).

The circumscription of the Verrucariaceae and the relationships among the main genera included in the family has been investigated by molecular methods (Gueidan et al. 2007, 2009; Savić et al. 2008). These studies showed that members of *Catapyrenium s.l.* were nested within different monophyletic groups. Gueidan et al. (2007, 2009) reported *Catapyrenium s. str.* as a sister group of *Placidiopsis* within the *Staurothele* group (=Lineage 4), a clade distantly related to the *Placidium* group nested within Lineage 3 and containing *Clavascidium*, *Neocatapyrenium*, *Placidium* and *Placopyrenium*. *Catapyrenium s. str.* and *Placidiopsis* formed a monophyletic group within the *Staurothele* group along with at least two putatively paraphyletic species of *Verrucaria*. Although both *Catapyrenium s. str.* and *Placidiopsis* seemed to be monophyletic, only two species of each genus were included in these analyses. Therefore the monophyly

of both genera is still considered an open question necessitating more taxon sampling to clarify whether these current genera are well delineated.

The main goals for this phylogenetic study were to evaluate whether *Catapyrenium s. str.* is a monophyletic group separate from *Placidiopsis* and whether the gain of transversally uniseptate ascospores is a synapomorphy for the latter genus. We also investigated whether *P. cinerascens* and *P. tenella* are conspecific.

#### MATERIALS AND METHODS

*Taxon sampling.*—To infer phylogenetic relationships of specimens selected for this study within family Verrucariaceae we conducted a first set of analyses based on the nuLSU locus (Verrucariaceae dataset, VD). In total VD included 114 sequences of Verrucariaceae, of which 81 were obtained from GenBank and 33 were obtained by the first author during this study. Two sequences from GenBank belonging to the Chaetothiales were used as outgroup.

For the second dataset (*Catapyrenium-Placidiopsis* dataset, C-PD) sequences of the nuITS and nuLSU rDNA were combined for a total of 26 specimens belonging to *Catapyrenium s. str.* and *Placidiopsis* genera. Two sequences of *Staurothele* were downloaded from GenBank and selected as outgroup based on previous phylogenies (Gueidan et al. 2007, 2009; Savić et al. 2008).

In summary a total of 59 sequences were produced in this study and 85 were downloaded from GenBank. Information on the material, area of collection, collector, location of voucher specimens and GenBank accession numbers is provided for newly obtained sequences (TABLE II). For sequences obtained from GenBank identification numbers are found after taxon names (FIGS. 1, 2). The classification of groups follows Gueidan et al. (2007) and taxon names follow Gueidan et al. (2009).

*Morphological study.*—Characters suitable for differentiating *Placidiopsis cinerascens* from *P. tenella* were identified in preliminary morphological and anatomical studies. Criteria for delimiting these two species followed Breuss (1996b). Longitudinal sections (14–16 µm thick) were obtained from thalli and perithecia with a freezing microtome. The sections were mounted and measured in water. Sixteen anatomical characters were analyzed in all specimens studied (thallus, epinecral layer, upper cortex, algal layer, medulla and lower cortex thickness, cell diameter in all layers, rhizohyphae, perithecium and exciple thickness and color, ascospore length and width). Ten measurements were made for each sample. All variables were normally distributed. We have provided results for the 12 characters with continuous variables (TABLE III).

We used generalized linear mixed models (GLMM; McCullagh and Nelder 1989, Breslow and Clayton 1993) that consider variation between specimens within groups. All specimens were separated into putative species groups based on the phylogenetic results (FIG. 2, groups A and B). We analyzed data with a two-level hierarchical approach

TABLE II. List of taxa for which DNA sequences were generated in this study (GenBank identification numbers of sequences obtained from GenBank are found after species names in FIG. 1), with countries and provinces/states of origin, collector names, herbarium names where specimens were deposited with herbarium accession numbers, and GenBank accession numbers

Species name	Origin	Collector	Voucher	GenBank accession number	
				nuITS	nuLSU
<i>Catapyrenium cinereum</i> 1 (Pers.) Körb.	Spain, Cantabria	G. Aragón, A. García & M. Prieto	MA 16299	GQ344599	GQ344586
<i>Catapyrenium cinereum</i> 2	Spain, Asturias	M. Prieto	MA 16301	GQ344598	GQ344587
<i>Catapyrenium dactylinum</i> Breuss	Mexico, Chihuahua	T.H. Nash	LI 412469	—	GQ344593
<i>Catapyrenium daedaleum</i> 1 (Kremp.) Stein	Spain, León	G. Amo & G. Aragón	MA 16297	GQ344596	GQ344582
<i>Catapyrenium daedaleum</i> 2	Spain, Cuenca	G. Aragón	MA 16296	GQ344597	GQ344583
<i>Catapyrenium exaratum</i> 1 Breuss	Argentina, Salta	G. Aragón & I. Martínez	BCRU 4920	GQ344600	GQ344589
<i>Catapyrenium exaratum</i> 2	Argentina, Salta	G. Aragón & I. Martínez	MA 16314	GQ344601	GQ344588
<i>Catapyrenium psoromoides</i> 1 (Borrer) R. Sant.	Spain, Toledo	G. Aragón & M. Prieto	MA 16298	GQ344594	GQ344584
<i>Catapyrenium psoromoides</i> 2	Portugal, Portagem	M.A.G. Otálora & M. Prieto	MA 16313	GQ344595	GQ344585
<i>Heteroplacidium acervatum</i> (Breuss) Breuss	Spain, Mallorca	O. Breuss	LI 271015	—	GQ344564
<i>Placidioopsis cinerascens</i> 1 (Nyl.) Breuss	Spain, Palencia	G. Aragón & M. Prieto	MA 16307	GQ344605	GQ344566
<i>Placidioopsis cinerascens</i> 2	Spain, Zaragoza	I. Martínez & M. Prieto	MA 16306	GQ344606	GQ344567
<i>Placidioopsis cinerascens</i> 3	Spain, Cáceres	M. Prieto	MA 16302	GQ344610	GQ344565
<i>Placidioopsis cinerascens</i> 4	Spain, Zaragoza	I. Martínez & M. Prieto	MA 16305	GQ344609	GQ344568
<i>Placidioopsis cinerascens</i> 5	Portugal, Alvados	M.A.G. Otálora & M. Prieto	MA 16309	GQ344607	GQ344569
<i>Placidioopsis cinerascens</i> 6	Spain, Mallorca	M. Prieto	MA 16304	GQ344614	GQ344571
<i>Placidioopsis cinerascens</i> 7	Spain, Madrid	M. Prieto	MA 16308	GQ344613	GQ344570
<i>Placidioopsis custnani</i> 1 (A. Massal.) Körb.	Spain, Huesca	M. Prieto	MA 16303	GQ344602	—
<i>Placidioopsis custnani</i> 2	Spain, La Rioja	I. Martínez & M. Prieto	MA 16310	GQ344604	GQ344578
<i>Placidioopsis custnani</i> 3	Portugal, Alvados	M.A.G. Otálora & M. Prieto	MA 16312	GQ344603	GQ344579
<i>Placidioopsis pseudocinerea</i> Breuss	Austria, Steiermark	O. Breuss & R. Türk	LI 281386	GQ344619	GQ344577
<i>Placidioopsis tenella</i> 1 (Nyl.) Zahlbr.	Spain, Castellón	M. Prieto	MA 16315	GQ344611	GQ344572
<i>Placidioopsis tenella</i> 2	Spain, Lérida	J. Pérez-Redondo	BCC 12680	GQ344608	GQ344574
<i>Placidioopsis tenella</i> 3	Turkey, Tekirdağ	A. Özdemir Türk	LI 281350	GQ344615	GQ344573
<i>Placidioopsis tenella</i> 4	Italy, Calabria	D. Puntillo	LI 281352	GQ344616	GQ344575
<i>Placidioopsis tenella</i> 5	Spain, Navarra	J. Etayo & O. Breuss	LI 136521	GQ344612	GQ344576
<i>Placidioopsis tiroliensis</i> 1 Breuss	Austria, Steiermark	E. Hörandl	LI 281351	GQ344617	GQ344580
<i>Placidioopsis tiroliensis</i> 2	Austria, Kärnten	R. Türk & O. Breuss	LI 281349	GQ344618	GQ344581
<i>Placidium pseudorufescens</i> Breuss	USA, New Mexico	R. D. Worthington	LI 566404	—	GQ344563
<i>Placidium tenellum</i> (Breuss) Breuss	Spain, Madrid	I. Martínez, M.A.G. Otálora & M. Prieto	MA 16300	—	GQ344562



TABLE II. Continued

Species name	Origin	Collector	Voucher	GenBank accession number	
				nuITS	nuLSU
<i>Placopyrenium caeruleopulvinum</i> 1 (Thomson) Breuss	USA, New Mexico	R. D. Worthington	COLO 465590	—	GQ344591
<i>Placopyrenium caeruleopulvinum</i> 2	USA, Arizona	T.H. Nash	DUKE 146691	—	GQ344592
<i>Placopyrenium trachyticum</i> (Hazsl.) Breuss	Belgium, Flemish Brabant	A. Aptroot	DUKE 139586	—	GQ344561
<i>Placopyrenium stanfordii</i> (Herre) K. Knudsen	USA, Arizona	K. Knudsen	LI 582724	—	GQ344590

with specimens nested within groups. All GLMM computations were performed with SAS Macro program GLIMMIX (Littel et al. 1996).

**DNA isolation and sequencing.**—Genomic DNA was obtained from fresh samples and herbarium specimens. Small samples were ground in liquid nitrogen. Total genomic DNA was extracted with DNeasy Plant Mini Kit (QIAGEN) according to manufacturer's instructions, with slight modifications described by Crespo et al. (2001). Dilutions (1 : 10 and 1 : 100) of genomic DNA were used for PCR amplifications of the nuclear nuITS and nuLSU rDNA regions. Fungal nuITS rDNA was amplified with primers ITS5 and ITS4 (White et al. 1990), while nuLSU was amplified with primers LR0R (Rehner and Samuels 1994) and LR7 (Vilgalys and Hester 1990). Because these primer pairs also amplified the algal component we designed the specific primers PRI1 and PRI2 (5'-GTGTGAAGCTCCTTCGAC-3' and 5'-AAAAATGGCCCACTAGTAACG-3'). We used combinations of these four primers for amplifications. Amplifications were performed in 25  $\mu$ L volumes containing a reaction mixture of 7.5  $\mu$ L QIAGEN Multiplex PCR Kit (HotStarTaq DNA Polymerase plus Multiplex PCR Buffer and dNTP mix), 2.5  $\mu$ L Q-Solution, 2.5  $\mu$ L primer mixture (10  $\mu$ M), 10.5  $\mu$ L dH<sub>2</sub>O and 2  $\mu$ L diluted genomic DNA. Amplifications were carried out in a PTC-100 Peltier thermal cycler and performed with these programs: initial denaturation at 94 C for 15 min for nuITS or 95 C for nuLSU, followed by 35 cycles of 94 C for nuITS or 95 C for nuLSU for 1 min, 54 C for 1 min for nuITS and 52 C for 0.40 min for nuLSU rDNA, 72 C for 1.3 min for nuITS or 3 min for nuLSU, followed by a final extension at 72 C for 10 min. PCR products were purified with the QIAquick PCR Purification Kit (QIAGEN) according to manufacturer's instructions. Purified PCR products were sequenced with the same amplification primers, as well as LR3R (Vilgalys and Hester 1990) for the nuLSU region. Both strands were sequenced with the ABI PRISM™ Dye Terminator Cycle Sequencing Ready reaction kit (Applied Biosystems) with these PCR settings: denaturation 3 min at 94 C, 25 cycles at 96 C for 10 s, 50 C for 5 s and 60 C for 4 min. Sequencing products were subjected to electrophoresis with a 3730 DNA analyser (Applied Biosystems).

**Sequence alignment.**—Sequences were subjected to BLAST

analyses for a first verification of their entities. They subsequently were assembled and edited with SeqNavigator 1.0.1 (Applied Biosystems) and aligned manually with McClade 4.06 (Maddison and Maddison 2003) with the help of the secondary structure of the nuLSU from *Saccharomyces cerevisiae* (Cannone et al. 2002) following the method described in Kjer (1995).

A total of 116 sequences were aligned for the VD matrix. Fourteen ambiguously aligned regions were recoded and subjected to specific step matrices obtained with program INAASE 2.3b (Lutzoni et al. 2000, <http://www.lutzonilab.net/downloads>), incorporating the phylogenetic signal from these regions without violating positional homology. Three ambiguous regions that were more than 100 bp long or highly variable (i.e. more than 32 character states) were recoded into 23 characters with the aid of program arc 1.5 (Kauff et al. 2003, Miadlikowska et al. 2003) with the nucleotide option as outlined by Reeb et al. (2004). Each of the 23 characters obtained with program arc were subjected to a specific weight: 1 for character 1, 0.25 for characters 2–5, 0.10 for characters 6–15 and 0.50 for characters 16–23. Introns were delimited manually and excluded from the phylogenetic analyses.

A total of 28 specimens were aligned for C-PD. Only one taxon with missing nuLSU sequences was included in the analysis (TABLE II). Ambiguous regions (sensu Lutzoni et al. 2000) and introns were delimited manually and excluded from the phylogenetic analyses. The sequences alignments have been deposited in TreeBASE (accession number SN4626).

**Phylogenetic analyses.**—We performed two weighted maximum parsimony (wMP) analyses: a first wMP analysis with VD was executed with nuLSU, including coded (INAASE and arc) characters. The second search with C-PD included both loci nuITS and nuLSU and exclusively unambiguously aligned sites.

wMP analyses were performed with PAUP\* 4.0b10 (Swofford 2002). In both analyses symmetric step matrices were created for unambiguous portions of each of the genomic regions as follows. The option “character status/full detail/hidden excluded characters” from the data menu in PAUP\* was implemented. From the resulting table the column states showing all nucleotide states and gaps found




FIG. 1. Phylogenetic relationships within the Verrucariaceae based on nuLSU data for 114 specimens of Verrucariaceae and two specimens selected as outgroup from the Chaetothoriales. Strict consensus tree of three equally most parsimonious trees generated with wMP using a backbone constraint tree derived from Guéidan et al. (2007). Sequences downloaded from GenBank are followed by their GenBank identification (GI) numbers. Specimens sequenced for this study are in boldface and

at each of the unambiguously aligned and nonconstant sites was saved as a separate text file. This file was used as an input for program StMatrix 2.1. (Lutzoni and Zoller, Duke University, <http://www.lutzonilab.net/downloads>), which generates a step matrix (in Nexus format) by calculating frequencies of reciprocal changes from one state to another and converting them into costs of changes with the negative natural logarithm of the frequencies (Felsenstein 1981, Wheeler 1990).

wMP analyses were performed with the heuristic search option with 1000 random addition sequences (RAS) replicates, TBR (tree bisection reconnection) branch-swapping and MULTREES option in effect. We performed a search in two steps for VD. In the first step we performed only 100 RAS (i.e. without branch swapping after the last sequence was added) while in the second step we searched for all equally parsimonious trees, saving all trees only when swapping a tree equal to or shorter than the shortest tree that we found in the first step.

All character states were treated as unordered, and gaps were considered as a fifth character. Nonparametric bootstrap (Felsenstein 1985) was used to assess statistical support of the clades with heuristic searches as described above on 1000 bootstrap datasets. Five RAS per bootstrap replicate were specified for VD and two for C-PD based on the efficiency of the searches based on the original datasets.

Analyses were performed three ways for VD bootstrap: saving no more than 10 trees per RAS in the first analysis, no more than 100 trees per RAS in the second and no more than 10 trees per RAS but with 10 000 bootstrap replicates in the third analysis. Bootstrap values were similar, consequently values from the first analysis are reported here (FIG. 1).

Preliminary analyses of VD failed to recover a well supported topology (results not shown). Based on multi-locus phylogenies of the Verrucariaceae by Gueidan et al. (2007), from which our initial nuLSU alignment was derived and to which we added nuLSU sequences generated for our study, we constructed a topological constraint tree for a subset of the deep nodes and including only sequences from Gueidan et al. (2007) (i.e. backbone constraint). This backbone constraint tree was constructed by hand with McClade 4.06 (Maddison and Maddison 2003). Internodes were included in this tree when they were highly supported (i.e.  $\geq 70\%$  bootstrap values and  $\geq 95\%$  posterior probabilities) at least in two of the three analysis based on weighted maximum parsimony (wMP), maximum likelihood bootstrap (ML BS) and Bayesian analysis (MB) in the three-gene tree of Gueidan et al. (2007). Specimens

sequenced for this study not included in the constraint (skeletal) tree were free to attach to any point on the trees during the search. Our approach did not assume any restrictions on the phylogenetic positioning of specimens sequenced for this study but enforced topological relationships between the remaining taxa only, as far as sufficiently supported in multigene phylogenies by Gueidan et al. (2007) that were based on more characters (i.e. nuLSU, nuSSU and *RPB1*).

Phylogenetic relationships and confidence for C-PD were inferred with weighted maximum parsimony, Bayesian analyses and bootstrap maximum likelihood on a combined nuITS-nuLSU dataset. The combinability of the datasets was assessed by separately comparing clades among trees based on the nuITS region and nuLSU (Mason-Gamer and Kellogg 1996). Because no conflict was detected it was assumed that the two datasets were congruent and could be combined.

The evolutionary models for Bayesian analyses were selected with the Akaike information criterion (AIC) as implemented in MrModeltest 2.2 (Nylander 2004). GTR+I+G was used for nuLSU, HKY+G for ITS 1 and ITS 2 and K80+I for ITS 5.8S in the C-PD. Datasets were analyzed with MrBayes 3.1.1 (Huelsenbeck and Ronquist 2001, Ronquist and Huelsenbeck 2003). Two analyses of four chains were run 5 000 000 generations starting from an initial random tree. A tree was saved every 100th generation. A burn-in sample of 150, 200 and 100 trees for nuITS, nuLSU and the combined dataset respectively were discarded. The remaining trees were used to estimate branch lengths with the SUMT command of MrBayes, and posterior probabilities (PP) were calculated with the majority rule consensus tree command in PAUP\*.

A maximum likelihood (ML) search for the most likely tree was completed on 1000 replicates with RAxML-VI-HPC (Stamatakis et al. 2005) with a GTRMIX model of molecular evolution. An additional constrained ML tree search (with the same backbone constraint tree mentioned above) was conducted on VD, and 1000 BS replicates were run for both datasets.

## RESULTS

*Verrucariaceae dataset.*—The data matrix included 1390 characters after exclusion of ambiguous regions and introns. Fourteen of the ambiguous regions were recovered by INAASE characters, and three of them by arc characters. Nine hundred four characters were excluded from the wMP analysis because they were

←

lack this number. Internodes with BS support from wMP and ML  $\geq 70\%$  are highlighted with thicker lines. Internodes with black lines were not part of the skeletal backbone constraint tree. Internodes with gray lines were part of the backbone constraint tree. Internodes with dashed lines were part of the backbone tree but not recovered with high support. The support values are ordered as wMP/ML BS. An asterisk over a bold branch indicates that this internode has a bootstrap support value of 100% for both wMP and ML. A dash indicates bootstrap support values  $< 50\%$ . Members of *Placopyrenium* are in gray boxes, showing the polyphyly of the genus. Dashed line box delimits *Catapyrenium s. str.* and *Placidioopsis* clades, which is the ingroup of the phylogenetic tree (FIG. 2). An arrow shows the position of *Catapyrenium dactylinum* outside *Catapyrenium s. str.*




FIG. 2. Phylogenetic relationships among 28 members of *Catapyrenium s. str.* and *Placidiopsis*, based on ML analysis of concatenated nuITS and nuLSU datasets. Two species of *Staurothele* were chosen as outgroup and are followed by their GenBank identification (GI) numbers. ML BS, PP and wMP BS are represented with the first, second and third numbers associated with branches. Branches in boldface indicate a support of PP  $\geq$  95%, and both ML and wMP BS  $\geq$  70%. An asterisk on a bold branch indicates that this internode has a value of 100% for all support estimates. A dash indicates bootstrap support



constant. Of the remaining sites, 338 were parsimony informative. Constant sites were included for maximum likelihood analyses.

The weighted maximum parsimony analysis with the backbone constraint tree resulted in three equal, most parsimonious trees, 3782.47 steps long, which were part of one island hit 663 times out of 1000 RAS. Resolution and support at deeper nodes largely reflect the backbone constraint in our analyses, but the presence of unconstrained taxa in their expected clades indicates that phylogenetic signal was not guided by our backbone constraint alone. This is the case for *Heteropladidium acervatum*, *Placidium pseudorufescens* and *Placidium tenellum* that appear in the *Placidium* group sensu Gueidan et al. (2007), together with other members of *Heteropladidium* and *Placidium* genera with high support values (FIG. 1).

All specimens of *Placidiopsis* and *Catapyrenium s. str.* (except *C. dactylinum*) sequenced by us, and which were free to attach anywhere on the tree, connected to the skeletal monophyletic quartet formed by *Placidiopsis cinerascens*, *P. custnani*, *Catapyrenium cinereum* and *C. daedaleum* from Gueidan et al. (2007). All new specimens we included connected to their respective genera thereby recovering the sister relation between these two genera (FIG. 1). In the clade containing specimens of *Catapyrenium s. str.* the internode part of the constrained backbone tree was recovered with high support value only with wMP (dashed line in FIG. 1). The core of *Catapyrenium s. str.* species are significantly supported by both analyses (82% wMP BS, 75% ML BS). This supported node includes all species of *Catapyrenium s. str.* except *C. dactylinum*. All *Placidiopsis* specimens included in this study formed a highly supported monophyletic group (100% wMP BS and ML BS). However interspecific relationships within this genus were not fully resolved. In the *Staurothele* group (sensu Gueidan et al. 2007) relationships not previously supported were highly supported in our analyses, such as the sister relationship between *Verrucaria caerulea* and *V. praetermissa* (wMP BS = 97%, ML BS = 96%) and the monophyletic group formed by these two species with *Catapyrenium s. str.*, *Placidiopsis* and *Staurothele* species (wMP BS = 77%, ML BS = 76%).

*Catapyrenium s. str.* was not recovered as monophyletic because *C. dactylinum* was found to be sister of *Placopyrenium caeruleopulvinum*. These two species, together with *Placopyrenium stanfordii*, belong to a

well supported clade that first split from the remaining members of the Verrucariaceae, including *Placocarpus schaeferi*, as well as the monophyletic genera *Verrucula* and *Wahlenbergiella*. These *Placopyrenium* species (*Placopyrenium* II) together with *Catapyrenium dactylinum*, *Placocarpus schaeferi* and the *Verrucula* group formed a clade that was highly supported by both analyses (100% BS support). The previously supported (Gueidan et al. 2007) and constrained relationship between *Placocarpus schaeferi* and the *Verrucula* clade was recovered but without significant support values (dashed line in FIG. 1).

The clade containing both species of *Placopyrenium* (*P. caeruleopulvinum* and *P. stanfordii*) and *Catapyrenium dactylinum* is only distantly related to the *Placopyrenium* group sensu Gueidan et al. (2007). This latter group comprises the type species of *Placopyrenium* (*P. bucekii*) and two *Verrucaria* species recently combined as *Placopyrenium* by Navarro-Rosinés et al. (2007) (i.e. *P. canellum* and *P. fuscellum*). Another species of *Placopyrenium*, *P. trachyticum*, not previously analyzed by Gueidan et al. (2007) was found to be closely related to *P. fuscellum* (FIG. 1). Despite results from Gueidan et al. (2007, 2009), which reported genus *Placopyrenium* as monophyletic, the addition of *Placopyrenium caeruleopulvinum* and *P. stanfordii* to their dataset resulted in *Placopyrenium s.l.* being polyphyletic. We refer to the clade containing the type species of *Placopyrenium*, as *Placopyrenium s. str.*, whereas we labeled the paraphyletic group containing the remaining species of *Placopyrenium* we analyzed as *Placopyrenium* II (FIG. 1).

Furthermore we obtained high bootstrap support for the *Placopyrenium* group (*Placopyrenium s. str.*) forming a monophyletic group with *Endocarpon* and *Polyblastia* groups sensu Gueidan et al. (2007) and for the clade composed of the *Placidium*, *Aquatic* and *Dermatocarpon* groups. The latter clade was found to be well supported as sister of *Baggioletta*. We must regard these relationships as tentative although well supported because only one ribosomal gene was used here, necessitating the use of a skeletal backbone constraint tree to resolve the phylogenetic placement of 16 additional species (representing five genera) within the Verrucariaceae tree of Gueidan et al. (2007).

*Catapyrenium-Placidiopsis dataset.*—The combined data matrix included 495 nuITS and 969 nuLSU

←

values < 50%. In the *Placidiopsis cinerascens-tenella* clade the presence/absence of involucrellum and geographical location of where the samples were obtained are shown after the name of each specimen.

TABLE III. Mean, standard deviation and range (in parentheses) of measurements (in micrometers) for *Placidiopsis cinerascens* and *P. tenella* specimens belonging to groups A and B, (FIG. 2)

Characters ( $\mu\text{m}$ )	Group A	Group B	significance
Thallus thickness	(150) $236.4 \pm 47.1$ (320)	(110) $213.9 \pm 48.9$ (300)	n.s.
Epinecral layer thickness	(3) $13.2 \pm 8.1$ (32.5)	(5) $18.8 \pm 10.9$ (50)	n.s.
Upper cortex thickness	(5) $20.8 \pm 8.6$ (37.5)	(5) $17.2 \pm 7.1$ (35)	n.s.
Upper cortex cells diameter	(4) $7 \pm 1.3$ (10)	(5) $7.3 \pm 1.6$ (11)	n.s.
Algal layer thickness	(62.5) $109.8 \pm 25.3$ (150)	(50) $101.6 \pm 34.42$ (175)	n.s.
Algal cells diameter	(3) $6.6 \pm 1.7$ (12)	(4) $6.5 \pm 1.8$ (11)	n.s.
Medullary cells diameter	(4) $7.7 \pm 1.7$ (11)	(5) $7.3 \pm 1.3$ (10)	n.s.
Rhizohyphae thickness	(3) $3.2 \pm 0.4$ (4)	(2.5) $3.2 \pm 0.5$ (4)	n.s.
Perithecium diameter	(140) $212 \pm 40.5$ (300)	(140) $191.7 \pm 30.9$ (250)	n.s.
Exciple thickness	(10) $17.2 \pm 4.9$ (32)	(10) $16 \pm 3.6$ (23)	n.s.
Ascospore length	(12) $16.6 \pm 1.6$ (21)	(12) $15.5 \pm 1.7$ (21)	n.s.
Ascospore width	(5.5) $6.1 \pm 0.38$ (7)	(5) $6.2 \pm 0.4$ (7)	n.s.

n.s. = no significant differences ( $P > 0.05$ ).

characters after exclusion of ambiguous regions and introns. From the 1464 unambiguously aligned sites, 1167 were excluded from the wMP analysis because they were constant. Of the remaining sites 227 were parsimony informative. Constant sites were included for the maximum likelihood and Bayesian analyses.

The weighted parsimony analysis of the combined dataset resulted in one equal, most parsimonious tree of 956.99 steps, which was part of one island hit 996 times out of 1000 RAS. The majority rule consensus tree of 99 800 sampled trees from Bayesian analysis was similar to the parsimony and maximum likelihood topologies, and no conflict was detected. The most likely tree from the ML analysis is presented (FIG. 2) with branch lengths and support values.

All analyses supported two sister monophyletic genera (100% PP, ML BS and wMP BS), consisting of *Placidiopsis* and *Catapyrenium* s. str. (excluding *C. dactylinum*). Species of *Placidiopsis* lacking rhizines, including *P. cinerascens*, *P. tenella*, *P. pseudocinerea* and *P. tirolensis*, formed a well supported monophyletic group sister of *P. custnani*, a species of *Placidiopsis* with a rhizine-like central holdfast.

A clade consisting of both *Placidiopsis cinerascens* and *P. tenella* was highly supported (99% ML BS, 100% PP and 96% wMP BS). These species were distinguished based on the presence or absence of an involucrellum (Breuss 1996b, Nimis and Martellos 2004) (FIG. 2). However the first author's observations led her to conclude that the presence of an involucrellum is not a reliable trait because it is not always associated with all ascomata in the same specimen or even in the same squamule of *P. tenella*. Therefore *Placidiopsis tenella* cannot be distinguished from *P. cinerascens* with this character.

Although the clade consisting of both *Placidiopsis cinerascens* and *P. tenella* seems to have

phylogenetic structure, showing a subclade with 100% support in all analyses, (group B in FIG. 2), and another subclade supported by ML and wMP BS (group A in FIG. 2), the pattern of presence/absence of an involucrellum for the specimens selected for this study does not correlate with the association of these specimens with groups A or B. A more detailed morphological and anatomical study of these two species confirms this result. No significant differences were found between groups A and B in the measured variables (TABLE III), and therefore no molecular or phenotypic traits support the maintenance of *Placidiopsis cinerascens* and *P. tenella* as distinct species. To find some differences between these two subclades we considered whether the specimens of each subclade showed some ecological or geographical trend, but members of both subclades live in the same ecological conditions (i.e. calciferous soils in dry, open habitats) and no geographical pattern seems to explain these two monophyletic groups (FIG. 2). Based on these results, *Placidiopsis tenella* is here reduced to synonymy with *Placidiopsis cinerascens*.

#### TAXONOMY

*Placidiopsis cinerascens* (Nyl.) Breuss, Plant Syst. Evol. 148:315. 1985 – Holotypus: Gallia merid., Beaucaille, W. Nylander (H-NYL 4021!)  
= *Placidiopsis tenella* (Nyl.) Zahlbr., Catal. Lich. Univ. I: 240. 1921 – Lectotypus designated by Cl. Roux in herbarium; Oran, Balansa (H-NYL 3944!)  
MycoBank MB 515120.

Note: We do not know the exact date of publication of *Endocarpon tenellum* in Annal Sci Nat (Nylander 1853b), however searching through some indirect

references we think that it could not have been published before the end of 1853. The publication of *Endocarpon cinerascens* in Bot Not (Nylander 1853a) is dated Oct–Nov 1853. With this information we decided to consider *Endocarpon cinerascens* as having priority over *E. tenellum*.

*Placidiopsis cinerascens* includes specimens with the upper surface greenish gray to brownish, with a variable epinecral layer and perithecia with or without apical involucrellum. The species is distributed in central Asia, Mediterranean Europe and Morocco, Mexico and SW North America in Mediterranean and arid climates.

Notes on the valid name of *Placidiopsis cartilaginea* (Nyl.) Vainio. Breuss (1996b) used the name *P. cartilaginea* in his revision of genus *Placidiopsis*, explaining the use of this name by Gueidan et al. in 2007. Even though *Verrucaria cinerea* var. *cartilaginea* was described in 1853 (i.e. before *Placidium custnani* A. Massal. [1856]), based on Article 11.2 of the ICBN this name has no priority due to the rank in which it was published. Therefore the accepted name is *Placidiopsis custnani* (A. Massal) Körber, Parerga Lich. 1963:305, which justifies the use of this name by Gueidan et al. in 2009.

#### DISCUSSION

The use of the nuLSU locus alone is clearly insufficient for resolving with high confidence relationships among a relatively large number of taxa representing species diversity across the Verrucariaceae. However it was sufficient to infer phylogenetic relationships of 16 species with regard to a core set of species for which relationships are fairly well established based on multilocus datasets. The efficiency and accuracy of tree searches are conditional on the quality of the backbone (skeletal) constraint trees used. In this study the backbone implemented was based on the most comprehensive phylogeny on Verrucariaceae conducted by Gueidan et al. (2007) with three loci (*RPB1*, nuLSU and nuSSU). We used this backbone tree, capturing known relationships within the Verrucariaceae, to first determine which taxa sequenced for this study were part of the *Staurothele* group, our main focus (FIG. 1). Based on this result we defined our ingroup, centered on *Catapyrenium s. str.* and *Placidiopsis*, for a more in-depth phylogenetic analysis based on nuITS and nuLSU, without the use of a backbone constraint tree.

The implementation of a backbone constraint tree let us take into consideration what is known based on

exhaustive multilocus phylogenies when conducting phylogenetic searches to infer relationships for additional taxa but on the basis of only one locus. Contrary to constraint trees that include all OTU of the dataset analyzed, internodes of backbone constraint trees will not necessarily be recovered in bootstrap analyses. For example the internode supporting the sister relationship of *Placocarpus schaeferi* and *Verrucula* received a bootstrap support value barely above 50% with wMP and below 50% with ML or the constrained relation between *Catapyrenium cinereum* and *C. daedaleum* was recovered only by wMP (FIG. 1).

This study demonstrates that *Catapyrenium s. str.* is not monophyletic because *C. dactylinum* is closely related to *Placopyrenium caeruleopulvinum*, *Placopyrenium stanfordii* and to a clade consisting of parasitic *Verrucula* species and *Placocarpus schaeferi*. This unexpected result could be explained by the presence of pycnidia on the specimen of *C. dactylinum*, being the only member of *Catapyrenium s. str.* with pycnidia. Other species of *Catapyrenium s. str.* lack pycnidia as do species of *Placidiopsis*. However members of *Verrucula* and *Placocarpus schaeferi*, along with *Catapyrenium dactylinum*, share *Dermatocarpon*-type pycnidia. Various authors have suggested that pycnidium type could be diagnostic for delimiting genera within the Verrucariaceae (Breuss 1990, Harada 1993, Gueidan et al. 2007). In addition *C. dactylinum* lacks a rhizohyphal weft (a constant feature in *Catapyrenium s. str.*), attaching to the substrate by its elongated basal ends. *Placocarpus schaeferi* (Breuss 1985), *Placopyrenium caeruleopulvinum* and *P. stanfordii* (Breuss 2002c) attach to the substrate by a central stipe. Both basal ends and stipes are rhizine-like structures originating from medullar tissue. Moreover all four taxa share a constant feature, the bluish-gray thallus produced by granular pruina. While looking for more shared characters among these species we realized that although *Catapyrenium dactylinum* is squamulose the squamules are aggregated, giving an areolate aspect to the thallus, similar to *Placopyrenium caeruleopulvinum*, which develops a crustose areolated thallus with convex to bullate areoles.

It is not surprising that *Placopyrenium caeruleopulvinum* and *P. stanfordii* occur in the same clade together with *Placocarpus schaeferi*. Hasse (1913) pointed out that *Placopyrenium stanfordii* (as *Dermatocarpon zalhbruckneri* Hasse) was morphologically similar to *Placocarpus schaeferi* (as *Dermatocarpon monstrosum* [Schaer.] Vain.) but differ in its uniform thallus and size of the ascospores. *P. caeruleopulvinum* and *P. stanfordii* are very similar, according to



Thomson (1987). Both species have a pale excipulum with only the mouth darkened, and the ascospores are larger in *Placopyrenium stanfordii*.

James et al. (2006) and Lawrey et al. (2007) have noted that genera within a given clade may have different life strategies. *Verrucula* species are parasitic, while *Placocarpus schaereri* has been reported to be parasitic only in the juvenile state (Zehetleitner 1978, Gueidan et al. 2009). However *P. caeruleopulvinum*, *P. stanfordii* and *Catapyrenium dactylinum* are not parasitic (Breuss 2000, 2002c, 2009). In our analysis the nonparasitic species formed a basal grade. The same evolutionary trend has been suggested to have taken place within the *Verruculopsis* group, according to Navarro-Rosinés et al. (2007).

Contrary to *Placopyrenium caeruleopulvinum* and *P. stanfordii*, *P. trachyticum* was found to belong to the *Placopyrenium* clade sensu Gueidan et al. (2007), here called *Placopyrenium s. str.* together with *P. buceki* and the recently combined *P. canellum* and *P. fuscillum* (Navarro-Rosinés et al. 2007). These members of *Placopyrenium* are characterized by having black-rimmed areoles or dark brown margins (Orange 2004, Breuss 2009). To date, *Placopyrenium* has been recognized as a monophyletic genus (Gueidan et al. 2007, 2009; Savić et al. 2008), but our results show the polyphyly of *Placopyrenium s.l.* A detailed study of this genus will be needed before proposing any taxonomical changes for species *P. caeruleopulvinum*, *P. stanfordii* and *C. dactylinum*.

Although the generic classification in family Verrucariaceae was based traditionally on ascospore septation (apart from structure of the thallus, presence or absence of hymenial alga, structure of the upper cortex and pycnidia; Zahlbruckner 1921–1922, Zschacke 1933–1934, Harada 1993, Breuss 1996a), recent molecular studies have shown that the ascospore type is often homoplasious and groups based on this character alone can result in polyphyly (Gueidan et al. 2007, 2009; Savić et al. 2008; Savić and Tibell 2009). Gueidan et al. (2007, 2009) questioned the monophyly of *Catapyrenium s. str.*, and *Placidiopsis* and suggested the possibility of combining these two genera into one genus. Results from our phylogenetic study support the recognition of *Catapyrenium s. str.* (excluding *C. dactylinum*) and *Placidiopsis* as distinct genera according to the traditional taxonomy of both genera (Breuss 1996a, b), with ascospore septation being a synapomorphic character for defining *Placidiopsis*.

In summary *Catapyrenium s. str.* is not monophyletic unless *C. dactylinum* is excluded from this genus and the same is true for *Placopyrenium* unless *P. stanfordii* and *P. caeruleopulvinum* are excluded, which we recommend for both genera but cannot

implement formally until the affiliation of these excluded species is determined through a detailed systematic study focusing on the *Verrucula-Placopyrenium-Wahlenbergiella* clade. Species delimitation has been problematic between *Placidiopsis cinerascens* and *P. tenella*. We have observed that the diagnostic involucrellum trait is present in specimens of both clades (FIG. 2), even though this character is not always present in all perithecia in the same thallus. The development of the involucrellum may vary and depends on the development of the thallus, which also depends on ecological conditions, such as exposure and humidity, according to Savić et al. (2008) and Thüs (2002). Because this character is not a constant feature we think it should not be employed for delimiting these two species. Both species were described by Nylander (1853a, b) who reported different ascospore lengths for these two species. In our study anatomical and morphological analyses have not shown differences between specimens of *Placidiopsis cinerascens* and *P. tenella* other than the expected within species level of variation. *Placidiopsis cinerascens* and *P. tenella* cannot be distinguished genotypically or phenotypically. Thus, *P. tenella* is reduced to synonymy with *P. cinerascens*.

#### ACKNOWLEDGMENTS

We are grateful for the aid of curators of BCN, COLO, DUKE, LI, NY and of C. Keller who loaned specimens, the critical comments of E. Gaya, B. Hodkinson and J. Miadlikowska, A. Escudero and R. García who provided statistical advice, R. Pérez who helped design PRI primers, A. Quintanar who helped with bibliographical search, O. Breuss for his useful comments and help and M. McMullen for the English revision. The authors also thank R. Belinchón, D. Martínez and J. Torres for help with fieldwork. This study was supported by research projects CGL2004-04795-C04-04, CGL2007-66066-C04-04 from the Spanish Ministry of Education and Science and a doctoral grant and support for two short stays in Natural History Museum Vienna and Duke University from Rey Juan Carlos University to M. Prieto.

#### LITERATURE CITED

- Breslow NE, Clayton DG. 1993. Approximate inference in generalized linear mixed models. *J Am Stat Assoc* 88:9–25.
- Breuss O. 1983. Eine neue *Placidiopsis*-Art (Lichenes, Verrucariaceae) aus Norwegen und den Alpen. *Plant Syst Evol* 142:247–250.
- . 1985. On the lichen genera *Placocarpus* and *Placidiopsis* (Verrucariaceae). *Plant Syst Evol* 148:313–315.
- . 1990. Die Flechtengattung *Catapyrenium* (Verrucariaceae) in Europa. *Stapfia* 23:1–153.



- . 1994. Die flechtengattungen *Catapyrenium* und *Placidiopsis* (Verrucariaceae) in Nordafrika. *Nova Hedwig* 58:229–237.
- . 1996a. Ein verfeinertes Gliederungskonzept für *Catapyrenium* (lichenisierte Ascomyceten, Verrucariaceae) mit einem Schlüssel für die bisher bekannten Arten. *Ann Naturhist Mus Wien* 98:35–50.
- . 1996b. Revision der Flechtengattung *Placidiopsis* (Verrucariaceae). *Öst Z Pilz* 5:65–94.
- . 2000. A peculiar new *Catapyrenium* species (lichenized Ascomycetes, Verrucariaceae) from Mexico. *Linzer Biol Beitr* 32:1053–1055.
- . 2001. *Catapyrenium*. In: McCarthy PM, ed. *Flora of Australia*. Vol. 58A, Lichens 3. Melbourne, Australia: ABRS/CSIRO. p 160–161.
- . 2002a. *Catapyrenium*. In: Nash TH III, Ryan BD, Gries C, Bungartz F, eds. *Lichen flora of the greater Sonoran Desert region I*. Tempe: Lichens Unlimited, Arizona State University. p 125–128.
- . 2002b. *Placidiopsis*. In: Nash TH III, Ryan BD, Gries C, Bungartz F, eds. *Lichen flora of the greater Sonoran Desert region I*. Tempe: Lichens Unlimited, Arizona State University. p 383–384.
- . 2002c. *Placopyrenium*. In: Nash TH III, Ryan BD, Gries C, Bungartz F, eds. *Lichen flora of the greater Sonoran Desert region I*. Tempe: Lichens Unlimited, Arizona State University. p 393–397.
- . 2009. A synopsis of the lichen genus *Placopyrenium* (Verrucariaceae), with descriptions of new taxa and key to all species. *Bibl Lichenol* 99:93–112.
- , Hansen ES. 1988. The lichen genera *Catapyrenium* and *Placidiopsis* in Greenland. *Plant Syst Evol* 159:95–105.
- Cannone JJ, Subramanian S, Schnare MN, Collett JR, D'Souza LM, Du Y, Feng B, Lin N, Madabusi LV, Muller KM, Pande N, Shang Z, Yu N, Gutell RR. 2002. The comparative RNA Website (CRW): an online database of comparative sequence and structure information for ribosomal, intron, and other RNA. *BMC Bioinformatics* 3:2–15.
- Crespo A, Blanco O, Hawksworth DL. 2001. The potential of mitochondrial DNA for establishing generic concepts in the parmelioid lichen. *Taxon* 50:807–819.
- Felsenstein J. 1981. A likelihood approach to character weighting and what it tells us about parsimony and compatibility. *Biol J Linn Soc* 16:183–196.
- . 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39:783–791.
- Gueidan C, Roux C, Lutzoni F. 2007. Using a multigene phylogenetic analysis to assess generic delineation and character evolution in Verrucariaceae (Verrucariales, Ascomycota). *Mycol Res* 111:1145–1168.
- , Savić S, Thüs H, Roux C, Keller C, Tibell L, Prieto M, Heiðmarsson S, Breuss O, Orange A, Fröberg L, Amtoft Wynns A, Navarro-Rosinés P, Krzewicka B, Pykälä J, Grube M, Lutzoni F. 2009. Generic classification of the Verrucariaceae (Ascomycota) based on molecular and morphological evidence: recent progress and remaining challenges. *Taxon* 58:184–208.
- Harada H. 1993. A taxonomic study on *Dermatocarpon* and its allied genera (Lichenes, Verrucariaceae) in Japan. *Nat Hist Res* 2:113–152.
- Harris RC. 1979. The genus *Placidiopsis* Beltr. (Lichenized Ascomycetes) new to North America as *Placidiopsis minor* sp. nov. *Michigan Bot* 18:57–58.
- Hasse HE. 1913. Additions to the lichen flora of southern California VIII. *Bryologist* 16:1–2.
- Hofstetter V, Miadlikowska J, Kauff F, Lutzoni F. 2007. Phylogenetic comparison of protein-coding versus ribosomal RNA-coding sequence data: a case study of the Lecanoromycetes (Ascomycota). *Mol Phylogenet Evol* 44:412–426.
- Huelsenbeck JP, Ronquist F. 2001. MrBayes: Bayesian inference of phylogenetic trees. *Bioinformatics* 17:754–755.
- Ihlen PG, Ekman S. 2002. Outline of phylogeny and character evolution in *Rhizocarpon* (Rhizocarpaceae, lichenized Ascomycota) based on nuclear ITS and mitochondrial SSU ribosomal DNA sequences. *Biol J Linn Soc* 77:535–546.
- James TY, Kauff F, Schoch C, Matheny B, Hofstetter V, Cox CJ, Celio G, Gueidan C, Fraker E, Miadlikowska J, Lumbsh T, Rauhut A, Reeb V, Arnold A, Amtoft A, Stajich JE, Hosaka K, Sung G, Johnson D, O'Rourke B, Crockett M, Binder M, Curtis JM, Slot JC, Wang Z, Wilson AW, Schueller A, Longcore JE, O'Donnell K, Mozley-Standridge S, Porter D, Letcher PM, Powell MJ, Taylor JW, White MM, Griffith GW, Davies DR, Humber RA, Morton JB, Sugiyama J, Rossmann AY, Rogers JD, Pfister DH, Hewitt D, Hansen K, Hambleton S, Shoemaker RA, Kohlmeyer J, Volkman-Kohlmeyer B, Spotts RA, Serdani M, Crous PW, Hughes KW, Matsuura K, Langer E, Langer G, Untereiner WA, Lücking R, Budel B, Geiser DM, Aptroot DM, Diederich P, Schmitt I, Schultz M, Yahr R, Hibbett DS, Lutzoni F, Mclaughlin DJ, Spatafora JW, Vilgalys R. 2006. Reconstructing the early evolution of Fungi using a six-gene phylogeny. *Nature* 443:818–822.
- Janex-Favre MC, Wagner J. 1986. Recherches ontogéniques et structurales sur les pycnides des Dermatocarpaceés (Pyrenolichens). *Bull Soc Mycol Fr* 102:161–182.
- Kauff F, Miadlikowska J, Lutzoni F. 2003. Arc: a program for ambiguous region coding. v 1.5. Durham, North Carolina: Duke University. <http://www.lutzonilab.net/downloads>
- Kjer KM. 1995. Use of rRNA secondary structure in phylogenetic studies to identify homologous positions: an example of alignment and data presentation from frogs. *Mol Phylogenet Evol* 4:314–330.
- Lawrey JD, Binder M, Diederich P, Molina MC, Sikaroodi M, Ertz D. 2007. Phylogenetic diversity of lichen-associated homobasidiomycetes. *Mol Phylogenet Evol* 44:778–789.
- Littell RC, Milliken GA, Stroup WW, Wolfinger RD. 1996. SAS system for linear models. Cary, North Carolina: SAS Institute Inc.
- Lutzoni F, Wagner P, Reeb V, Zoller S. 2000. Integrating ambiguously aligned regions of DNA sequences in phylogenetic analyses without violating positional homology. *Syst Biol* 49:628–651.
- Maddison WP, Maddison DR. 2003. MacClade: analysis of

- phylogeny and character evolution. version 4.6. Sunderland, Massachusetts: Sinauer Associates.
- Mason-Gamer RJ, Kellogg EA. 1996. Testing for phylogenetic conflict among molecular datasets in the tribe Triticeae (Gramineae). *Syst Biol* 49:524–545.
- McCullagh P, Nelder JA. 1989. Generalized linear models. London: Chapman & Hall.
- Ménard T, Roux C. 1995. *Placopyrenium bucekii* et remarques sur les Verrucariaceae (Verrucariales, Lichenes). *Mycotaxon* 53:129–159.
- Miadlikowska J, Lutzoni F, Goward T, Zoller S, Posada D. 2003. New approach to an old problem: incorporating signal from gap-rich regions of ITS and rDNA large subunit into phylogenetic analyses to resolve the *Peltigera canina* species complex. *Mycologia* 95:1181–1203.
- Miller AN, Huhndorf SM. 2004. A natural classification of *Lasiosphaeria* based on nuclear LSU rDNA sequences. *Mycol Res* 108:26–34.
- Naesborg RR, Ekman S, Tibell L. 2007. Molecular phylogeny of the genus *Lecania* (Ramalinaceae, lichenized Ascomycota). *Mycol Res* 111:581–91.
- Navarro-Rosinés P, Roux C, Gueidan C. 2007. La Genroj *Verrucula* kaj *Verruculopsis* (Verrucariaceae, Verrucariales). *Bull Soc Linn Provence* 58:133–180.
- Nimis PL, Martellos S. 2004. Keys to the lichens of Italy I. Terricolous species. Trieste: Edizioni Goliardiche. 341 p.
- Nylander JA. 2004. MrModeltest version 2. Program distributed by the author. Uppsala University Evolutionary Biology Centre.
- Nylander W. 1853a. Collectanea lichenologica in Gallia meridionali et Pyrenaeis. *Bot Not* 10–11:151–165.
- . 1853b. Lichenes Algeriensis novi. *Annal Sci Nat Bot* 20(3):315–320.
- Orange A. 2004. The *Verrucaria fuscella* group in Great Britain and Ireland. *Lichenologist* 36:173–182.
- Otálora MAG, Aragón G, Molina MC, Martínez I, Lutzoni F. 2010. Disentangling the *Collema-Leptogium* complex through a molecular phylogenetic study of the Collemataceae (Peltigerales, lichen-forming Ascomycota). *Mycologia* 102:279–290.
- Prieto M, Aragón G, Martínez I, Breuss O. 2008. A new species of *Anthracoarpon* (Verrucariaceae) from Argentina. *Bryologist* 111:128–132.
- Reeb V, Lutzoni F, Roux C. 2004. Contribution of *RPB2* to multilocus phylogenetic studies of the Pezizomycotina (Euscomycetes, Fungi) with special emphasis on the lichen-forming Acarosporaceae and evolution of poly-spory. *Mol Phylogenet Evol* 32:1036–1060.
- Rehner SA, Samuels GJ. 1994. Taxonomy and phylogeny of *Gliocladium* analysed from nuclear large subunit ribosomal DNA sequences. *Mycol Res* 98:625–634.
- Ronquist F, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19:1572–1574.
- Roux C. 2005. Likenoj de Okcidenta Eŭropo. Suplemento 4a: eltiraĵoj (extraits). *Bull Inform Assoc Fr Lichénol* 30:5–14.
- Savić S, Tibell L. 2009. Taxonomy and species delimitation in *Sporodictyon* (Verrucariaceae, Verrucariales) in northern Europe and the adjacent Arctic—reconciling molecular and morphological data. *Taxon* 58:585–605.
- , ———, Gueidan C, Lutzoni F. 2008. Molecular phylogeny and systematics of *Polyblastia* (Verrucariaceae, Eurotiomycetes) and allied genera. *Mycol Res* 112:1307–1318.
- Staiger B. 2002. Die Flechtenfamilie *Graphidaceae*. Studien in Richtung einer natürlicheren Gliederung. *Bibl Lichenol* 85:1–526.
- Stamatakis A, Ludwig T, Meier H. 2005. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22:2688–2690.
- Swofford D. 2002. PAUP\*: phylogenetic analysis using parsimony (\*and other methods). Version 4. Sunderland, Massachusetts: Sinauer Associates.
- Thomson JW. 1987. The lichen genera *Catapyrenium* and *Placidopsis* in North America. *Bryologist* 90:27–39.
- Thüs H. 2002. Taxonomie, Verbreitung und Ökologie silicoler Süßwasserflechten im außeralpinen Mitteleuropa. *Bibl Lichenol* 83:1–214.
- Vilgalys R, Hester M. 1990. Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *J Bacteriol* 172:4238–4246.
- Wheeler WC. 1990. Combinatorial weights in phylogenetic analysis: a statistical parsimony procedure. *Cladistics* 6: 269–275.
- White TJ, Bruns T, Lee S, Taylor JW. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, eds. PCR protocols: a guide to methods and applications. San Diego: Academic Press. p 315–322.
- Zahlbruckner A. 1921–1922. *Catalogus Lichenum Universalis* Band 1. Leipzig: Gebrüder Borntraeger.
- Zehetleitner G. 1978. Über einige parasitische Arten der Flechtengattung *Verrucaria*. *Nova Hedwig* 29:683–734.
- Zschacke H. 1933–1934. Epigloeaceae, Verrucariaceae und Dermatocarpaceae. In: Zahlbruckner A, ed. Dr L. Rabenhorst's Kryptogamen-Flora von Deutschland, Österreich und der Schweiz, 2. Aufl., Bd. 9, Abt. 1, Teil 1. Leipzig: Akademische Verlagsanstalt. p 44–695.